

COMPLEX PRODUCT STRUCTURES ON LIE ALGEBRAS

ADRIÁN ANDRADA AND SIMÓN SALAMON

Abstract. A study is made of real Lie algebras admitting compatible complex and product structures, including numerous 4-dimensional examples. If $\mathfrak{g}$ is a Lie algebra with such a structure then its complexification has a hypercomplex structure. It is shown in addition that $\mathfrak{g}$ splits into the sum of two left-symmetric subalgebras. Interpretations of these results are obtained that are relevant to the theory of both hypercomplex and hypersymplectic manifolds and their associated connections.

MSC. 17B60; 53C15, 53C30

1. INTRODUCTION

A complex structure on a real Lie algebra $\mathfrak{g}$ is an endomorphism J of $\mathfrak{g}$ satisfying $J^2 = -\mathbf{1}$ and the usual integrability condition expressed in terms of Lie brackets. Let $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ denote the complexification of $\mathfrak{g}$, and $\sigma : \mathfrak{g}^{\mathbb{C}} \rightarrow \mathfrak{g}^{\mathbb{C}}$ the corresponding conjugation. The integrability of J is then equivalent to a splitting

$$\mathfrak{g}^{\mathbb{C}} = \mathfrak{g}^{1,0} \oplus \mathfrak{g}^{0,1},$$

where $\mathfrak{g}^{1,0}$ and $\mathfrak{g}^{0,1}$ are complex Lie subalgebras of $\mathfrak{g}^{\mathbb{C}}$ and $\mathfrak{g}^{0,1} = \sigma(\mathfrak{g}^{1,0})$. A product structure on $\mathfrak{g}$ is an endomorphism E of $\mathfrak{g}$ which satisfies $E^2 = \mathbf{1}$ and an even simpler integrability condition. Let

$$\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$$

be the splitting in which $\mathfrak{g}_{\pm}$ denotes the eigenspaces corresponding to the eigenvalue ± 1 of E . Then the integrability of E is equivalent to requiring that $\mathfrak{g}_+$, $\mathfrak{g}_-$ are both Lie subalgebras of $\mathfrak{g}$.

In this article we are interested in another structure on $\mathfrak{g}$ which arises from the combination of the two above. Namely, a *complex product structure* on a real Lie algebra $\mathfrak{g}$ is a pair $\{J, E\}$ where J is a complex structure on $\mathfrak{g}$, E is a product structure on $\mathfrak{g}$, and $JE = -EJ$. In this case, the subalgebras $\mathfrak{g}_{\pm}$ satisfy $\mathfrak{g}_- = J\mathfrak{g}_+$. We summarize the basic definitions in §2, and provide some simple examples and constructions of such structures. Whilst the concept of a product structure is a relatively elementary one, we shall see that complex product structures are altogether more subtle and their classification far from easy.

A complex product structure on a Lie algebra is an analogue of a hypercomplex structure, i.e. a pair of anticommuting complex structures. Indeed, a complex product structure is a ‘latent hypercomplex structure’ in the sense that a complex product structure on a Lie algebra $\mathfrak{g}$ determines a hypercomplex structure on the real Lie algebra $(\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$ underlying $\mathfrak{g}^{\mathbb{C}}$. This result, that we establish in §3, provided our original interest in the subject. We also

explain how a standard complex product structure on $\mathfrak{gl}(2n, \mathbb{R})$ gives rise to a hypercomplex structure on $\mathfrak{gl}(2n, \mathbb{C})$. In general, $(\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$ also admits a complex product structure, so the process can be continued so as to obtain an infinite family of hypercomplex structures.

Complex product structures are closely related to a number of other algebraic concepts, and these relationships are pursued in §4. If $(\mathfrak{g}, J, E)$ is a Lie algebra with a complex product structure as above, then $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ is an example of a double Lie algebra [21]. Actually, following [22], $(\mathfrak{g}_+, \mathfrak{g}_-)$ is a matched pair of Lie algebras for certain representations $\rho : \mathfrak{g}_+ \rightarrow \mathfrak{gl}(\mathfrak{g}_-)$ and $\mu : \mathfrak{g}_- \rightarrow \mathfrak{gl}(\mathfrak{g}_+)$. The isomorphism $J : \mathfrak{g}_+ \rightarrow \mathfrak{g}_-$ then allows us to construct left-symmetric algebra (LSA) structures on $\mathfrak{g}_+$ and $\mathfrak{g}_-$. This implies that neither $\mathfrak{g}_+$ nor $\mathfrak{g}_-$ are semisimple [10], though we do not know whether there are examples with $\mathfrak{g}$ semisimple. Theorems 4.1 and 4.2 provide a complete characterization of Lie algebras carrying complex product structures in terms of pairs of LSA structures, and we also examine the special situation in which $\mathfrak{g}$ is itself an LSA.

Whilst the focus of this paper is algebraic, the correspondence between flat torsion-free connections on a Lie group and LSA structures on its Lie algebra allows us to give various geometrical interpretations of the theory developed thus far. This we do in §5. Suppose that $\mathfrak{g}$ has a complex product structure, and is the Lie algebra of a Lie group G . The endomorphisms J, E can be regarded as left-invariant tensors on G still satisfying $JE = -EJ$. This makes G a *complex product manifold*, a concept that is most easily defined in terms of a $GL(n, \mathbb{R})$ -structure for a diagonal inclusion $GL(n, \mathbb{R}) \subset GL(2n, \mathbb{R})$. We explain how properties of this type of manifold reflect the theory of §4, referring the reader to [1] for a more complete treatment.

A complex product manifold M possesses two involutive totally real distributions $T_{\pm}$. It also has a unique torsion-free connection ∇^{CP} for which J, E are parallel, and the integral submanifolds of $T_{\pm}$ are flat with respect to ∇^{CP} . Thus, complex product manifolds are integrable in a rather strong sense. When $M = G$ is a Lie group, we obtain a local double Lie group (G, G_+, G_-) in which each of the subgroups $G_{\pm}$ has a flat affine structure. We also show that ∇^{CP} gives rise to the Obata connection of the associated hypercomplex structure on $G^{\mathbb{C}}$; indeed the two connections share many properties.

There are natural links between complex product structures and symplectic geometry, since a class of left-symmetric algebras are those arising from Lie groups with a left-invariant symplectic form [8]. It therefore makes sense to look for complex product structures on Lie algebras for which $\mathfrak{g}_+, \mathfrak{g}_-$ are both *symplectic* Lie algebras. Algebraically, this means that $n = 2m$ is even and the structure group $GL(n, \mathbb{R})$ reduces to $Sp(m, \mathbb{R})$. Geometrically, this arises when an associated Lie group is endowed with a *hypersymplectic* structure in the sense of [17]. We explain this briefly in §5, but do not pursue this aspect of the theory in the present paper.

In §6, we consider examples of 4-dimensional Lie algebras carrying complex product structures, referring in part to the classification of complex structures given by Ovando [26] and Snow [30]. We determine the associated affine Lie groups and the hypercomplex structures on their complexifications. We should emphasize though that the complex product structures considered in this article can exist in dimensions $2n \not\equiv 0 \pmod{4}$, and we expect a subsequent analysis of the case $n = 3$ to prove fruitful.

Acknowledgments. The first author was supported by grants from CONICET, FONCYT and SECYT-UNC (Argentina). The authors are grateful to I. Dotti, A. Fino and R. Miatello for various valuable suggestions, and to N. Andruskiewitsch and D. Burde for help with the references.

2. PRELIMINARIES AND BASIC RESULTS

We begin by recalling some definitions which will be used throughout this work. All Lie algebras will be finite dimensional and defined over $\mathbb{R}$, unless explicitly stated.

An *almost complex structure* on a Lie algebra $\mathfrak{g}$ is a linear endomorphism $J : \mathfrak{g} \longrightarrow \mathfrak{g}$ satisfying $J^2 = -\mathbf{1}$. If J satisfies the condition

$$(1) \quad J[X, Y] = [JX, Y] + [X, JY] + J[JX, JY] \quad \text{for all } X, Y \in \mathfrak{g},$$

we will say that J is *integrable* and we will call it a *complex structure* on $\mathfrak{g}$. Note that the dimension of a Lie algebra carrying an almost complex structure must be even. Given an almost complex structure J on $\mathfrak{g}$, there is a splitting of $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ into the sum of two subspaces

$$(2) \quad \mathfrak{g}^{\mathbb{C}} = \mathfrak{g}^{1,0} \oplus \mathfrak{g}^{0,1},$$

where $\mathfrak{g}^{1,0}$ and $\mathfrak{g}^{0,1}$ are the eigenspaces corresponding to the eigenvalues $\pm i$ of the complexification of J . It is easy to see that J is integrable if and only if both $\mathfrak{g}^{1,0}$ and $\mathfrak{g}^{0,1}$ are complex subalgebras of $\mathfrak{g}^{\mathbb{C}}$.

The splitting (2) induces a decomposition of $(\mathfrak{g}^{\mathbb{C}})^*$ as

$$(\mathfrak{g}^{\mathbb{C}})^* = (\mathfrak{g}^{1,0})^* \oplus (\mathfrak{g}^{0,1})^*$$

and then

$$\bigwedge^k (\mathfrak{g}^{\mathbb{C}})^* = \bigoplus_{p+q=k} \Lambda^{p,q},$$

where $\Lambda^{p,q} \cong \bigwedge^p (\mathfrak{g}^{1,0})^* \otimes \bigwedge^q (\mathfrak{g}^{0,1})^*$ is the space of (p, q) -forms relative to J .

Recall that the Lie bracket in $\mathfrak{g}$ can be thought of as a linear map $[\cdot, \cdot] : \bigwedge^2 \mathfrak{g} \longrightarrow \mathfrak{g}$. In this way, we can consider its transpose $d : \mathfrak{g}^* \longrightarrow \bigwedge^2 \mathfrak{g}^*$, defined as follows:

$$(df)(X \wedge Y) = -f([X, Y]) \quad \text{for all } f \in \mathfrak{g}^*, X, Y \in \mathfrak{g}.$$

It is well known that J is integrable if and only if

$$d(\Lambda^{1,0}) \subseteq \Lambda^{2,0} \oplus \Lambda^{1,1}.$$

Whilst an almost complex structure on a Lie algebra $\mathfrak{g}$ makes the latter a complex vector space, $\mathfrak{g}$ need not be a complex Lie algebra since the Lie bracket might not be $\mathbb{C}$ -bilinear. This problem may be overcome in the following situation. Let $\mathfrak{g}$ be a Lie algebra with an almost complex structure I which satisfies

$$(3) \quad [IX, Y] = I[X, Y] \quad \text{for all } X, Y \in \mathfrak{g},$$

It is straightforward to verify that I is integrable, since (3) is stronger than (1). Then $\mathfrak{g}$ can be viewed as a complex Lie algebra, with multiplication by i given by the endomorphism I , since (3) implies the $\mathbb{C}$ -bilinearity of the Lie bracket.

Condition (3) is equivalent to $I \operatorname{ad}(X) = \operatorname{ad}(X)I$ for all $X \in \mathfrak{g}$, i.e. $\operatorname{ad}(X)$ is a complex transformation on $\mathfrak{g}$, and it is also equivalent to $d(\Lambda^{1,0}) \subseteq \Lambda^{2,0}$. Conversely, if $\mathfrak{h}$ is a complex Lie algebra, let $\mathfrak{h}_{\mathbb{R}}$ denote the underlying real Lie algebra. Then the endomorphism I of $\mathfrak{h}_{\mathbb{R}}$ given by multiplication by i , i.e. $IX = iX$ for all $X \in \mathfrak{h}_{\mathbb{R}}$, defines a complex structure on $\mathfrak{h}_{\mathbb{R}}$ satisfying (3).

Next, we define another kind of structure on a Lie algebra which is analogous to a complex structure. An *almost product structure* on $\mathfrak{g}$ is a linear endomorphism $E : \mathfrak{g} \longrightarrow \mathfrak{g}$ satisfying $E^2 = \mathbf{1}$ (and not equal to $\pm \mathbf{1}$). It is said to be *integrable* if

$$(4) \quad E[X, Y] = [EX, Y] + [X, EY] - E[EX, EY] \quad \text{for all } X, Y \in \mathfrak{g}.$$

An integrable almost product structure will be called a *product structure*. If $\dim \mathfrak{g}_+ = \dim \mathfrak{g}_-$, where $\mathfrak{g}_\pm$ is the eigenspace of $\mathfrak{g}$ associated to the eigenvalue ± 1 of E , then the product structure E is called a *paracomplex structure* [19, 20]. In this case, $\mathfrak{g}$ also has even dimension.

Given an almost product structure E on $\mathfrak{g}$, we have a decomposition $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$ into the direct sum of two linear subspaces, the eigenspaces associated to E , which induces a splitting $\mathfrak{g}^* = A_+ \oplus A_-$ and then

$$\bigwedge^k \mathfrak{g}^* = \bigoplus_{p+q=k} A^{p,q},$$

where $A^{p,q} \cong \bigwedge^p A_+ \otimes \bigwedge^q A_-$. It is easy to verify that E is integrable if and only if $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are both Lie subalgebras of $\mathfrak{g}$, and this is in turn equivalent to

$$(5) \quad d(A^{1,0}) \subseteq A^{2,0} \oplus A^{1,1}, \quad d(A^{0,1}) \subseteq A^{0,2} \oplus A^{1,1}.$$

Verification of these statements can be found within the proof of Proposition 2.2.

Lie algebras carrying either a complex structure or a product structure have many interesting properties that have been exhaustively studied. Our interest lies in an appropriate combination of these two structures:

Definition 2.1. A *complex product structure* on the Lie algebra $\mathfrak{g}$ is a pair $\{J, E\}$ of a complex structure J and a product structure E satisfying $JE = -EJ$.

The condition $JE = -EJ$ implies that the eigenspaces corresponding to the eigenvalues $+1$ and -1 of E have the same dimension, showing that E is in fact a paracomplex structure on $\mathfrak{g}$.

The endomorphism $F := JE$ satisfies $F^2 = \mathbf{1}$, and overall $\{J, E, F\}$ obey the rules

$$(6) \quad \begin{aligned} -J^2 &= E^2 = F^2 = \mathbf{1}, \\ JE &= F, \quad EF = -J, \quad FJ = E, \end{aligned}$$

satisfied by the so-called paraquaternionic numbers [15]. It is easy to verify that (4) is satisfied by F in place of E . Indeed $\mathfrak{g}$ has a circle's worth $\{\cos_\theta E + \sin_\theta F\}$ of product structures, and a corresponding ‘pencil’ of subalgebras $\mathfrak{g}_\theta$ with $\mathfrak{g}_0 = \mathfrak{g}_+$,

$$\mathfrak{g}_{\pm\pi/2} = \{X \pm JX : X \in \mathfrak{g}_+\},$$

and $\mathfrak{g}_\pi = \mathfrak{g}_-$.

Proposition 2.2. *Let $\mathfrak{g}$ be a Lie algebra. The following statements are equivalent:*

- (i) $\mathfrak{g}$ has a complex product structure,
- (ii) $\mathfrak{g}$ has a complex structure J and can be decomposed as $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$, where $\mathfrak{g}_+, \mathfrak{g}_-$ are Lie subalgebras of $\mathfrak{g}$ and $\mathfrak{g}_- = J\mathfrak{g}_+$.
- (iii) $\mathfrak{g}$ has a complex structure J and $\mathfrak{g}^*$ can be decomposed as $\mathfrak{g}^* = A_+ \oplus A_-$, where A_+, A_- are subspaces satisfying (5) and $A_- = JA_+$.

Proof. (i) $\iff$ (ii). If $\mathfrak{g}$ has a complex product structure $\{J, E\}$, let $\mathfrak{g}_\pm$ denote the eigenspace corresponding to the eigenvalue ± 1 of E . The integrability of E implies that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are Lie subalgebras of $\mathfrak{g}$ and $JE = -EJ$ implies $\mathfrak{g}_- = J\mathfrak{g}_+$.

Conversely, if (ii) holds, set $E|_{\mathfrak{g}_+} = \mathbf{1}$ and $E|_{\mathfrak{g}_-} = -\mathbf{1}$. Then it is easy to verify that $\{J, E\}$ determine a complex product structure on $\mathfrak{g}$.

(ii) $\iff$ (iii). Assuming (ii), let $A_+ \subset \mathfrak{g}^*$ (respectively, $A_- \subset \mathfrak{g}^*$) denote the annihilator of $\mathfrak{g}_-$ (respectively, $\mathfrak{g}_+$). Since $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are subalgebras of $\mathfrak{g}$, one immediately checks (5). The complex structure J induces a complex structure J on the vector space $\mathfrak{g}^*$ by setting $Jf = f \circ J$. Then $\mathfrak{g}_- = J\mathfrak{g}_+$ implies $A_- = JA_+$.

Conversely, if (iii) holds, set $\mathfrak{g}_+ = \{X \in \mathfrak{g} : f(X) = 0 \text{ for all } f \in A_-\}$ and $\mathfrak{g}_- = \{X \in \mathfrak{g} : f(X) = 0 \text{ for all } f \in A_+\}$. Conditions (5) imply at once that $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are Lie subalgebras of $\mathfrak{g}$ and $\mathfrak{g}_- = J\mathfrak{g}_+$ follows from $A_- = JA_+$. $\square$

Definition 2.3. Let $(\mathfrak{g}, J, E)$ and $(\mathfrak{g}', J', E')$ be two Lie algebras with complex product structures. We shall say that these complex product structures are *equivalent* if there exists an isomorphism of Lie algebras $\phi : \mathfrak{g} \longrightarrow \mathfrak{g}'$ such that $\phi J = J' \phi$ and $\phi E = E' \phi$.

The equivalence of Lie algebras with complex product structures can be reformulated in the following manner: according to (ii) of Proposition 2.2, we have a decomposition into a sum of subalgebras $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$ and $\mathfrak{g}' = \mathfrak{g}'_+ \oplus \mathfrak{g}'_-$ with $\mathfrak{g}_- = J\mathfrak{g}_+$, $\mathfrak{g}'_- = J'\mathfrak{g}'_+$. Then $\phi E = E' \phi$ means that $\phi(\mathfrak{g}_\pm) = \mathfrak{g}'_\pm$, i.e. ϕ maps each eigenspace for E onto the corresponding eigenspace for E' . Also $\phi J = J' \phi$ means that the natural extension of ϕ to an isomorphism $\phi^{\mathbb{C}} : \mathfrak{g}^{\mathbb{C}} \longrightarrow (\mathfrak{g}')^{\mathbb{C}}$ satisfies $\phi^{\mathbb{C}}(\mathfrak{g}^{1,0}) = (\mathfrak{g}')^{1,0}$ and $\phi^{\mathbb{C}}(\mathfrak{g}^{0,1}) = (\mathfrak{g}')^{0,1}$, where the splittings $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g}^{1,0} \oplus \mathfrak{g}^{0,1}$, $(\mathfrak{g}')^{\mathbb{C}} = (\mathfrak{g}')^{1,0} \oplus (\mathfrak{g}')^{0,1}$ are as in (2).

A definition that will be useful for our purposes is the following, which appeared in [21].

Definition 2.4. Three Lie algebras $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ form a *double Lie algebra* if $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are Lie subalgebras of $\mathfrak{g}$ and $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$ as vector spaces.

Observe that a double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ gives a product structure $E : \mathfrak{g} \longrightarrow \mathfrak{g}$ on $\mathfrak{g}$, where $E|_{\mathfrak{g}_+} = \mathbf{1}$ and $E|_{\mathfrak{g}_-} = -\mathbf{1}$. Conversely, a product structure on the Lie algebra $\mathfrak{g}$ gives rise to a double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$, where $\mathfrak{g}_\pm$ is the eigenspace associated to the eigenvalue ± 1 of E . Thus, a complex product structure $\{J, E\}$ on $\mathfrak{g}$ gives rise to a double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ with $\mathfrak{g}_- = J\mathfrak{g}_+$. The complex structure J determines a vector space isomorphism $J : \mathfrak{g}_+ \longrightarrow \mathfrak{g}_-$ which satisfies the condition (1) on $\mathfrak{g}$. In particular, $\dim \mathfrak{g}_+ = \dim \mathfrak{g}_-$.

Let us assume now we begin with a double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ such that $\dim \mathfrak{g}_+ = \dim \mathfrak{g}_-$. Consider a linear isomorphism $\varphi : \mathfrak{g}_+ \longrightarrow \mathfrak{g}_-$ and define an endomorphism J of $\mathfrak{g}$ by $J(X + A) = -\varphi^{-1}(A) + \varphi(X)$ for $X \in \mathfrak{g}_+$ and $A \in \mathfrak{g}_-$. J is clearly an almost complex structure on $\mathfrak{g}$ but, if we want J to be integrable, we need φ to satisfy the extra condition

$$(7) \quad \varphi[X, Y] + \varphi^{-1}[\varphi X, \varphi Y] = [\varphi X, Y] + [X, \varphi Y], \quad X, Y \in \mathfrak{g}_+.$$

These observations are summarized by

Proposition 2.5. *A double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ with $\dim \mathfrak{g}_+ = \dim \mathfrak{g}_-$ is associated to a complex product structure on the Lie algebra $\mathfrak{g}$ if and only if there exists a linear isomorphism $\varphi : \mathfrak{g}_+ \longrightarrow \mathfrak{g}_-$ such that (7) holds.*

We now wish to exhibit some simple examples of Lie algebras carrying complex product structures. A natural class to begin with is the class of real Lie algebras underlying a complex Lie algebra, since we have already mentioned that they are naturally endowed with a complex structure. Nevertheless, it is impossible to find interesting examples of complex product structures among these algebras, due to

Proposition 2.6. *Suppose that $\mathfrak{g}$ admits a complex product structure given by a complex structure I and a product structure E where I satisfies (3). Then $\mathfrak{g}$ is abelian.*

Proof. There is a splitting $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$, with $\mathfrak{g}_+$ and $\mathfrak{g}_- = I\mathfrak{g}_+$ Lie subalgebras of $\mathfrak{g}$. If $U, V \in \mathfrak{g}_+$ then

$$E[U, IV] = EI[U, V] = -IE[U, V] = -I[U, V] = -[U, IV],$$

which shows $[U, IV] \in \mathfrak{g}_-$, and

$$E[U, IV] = -E[I(IU), IV] = -EI[IU, IV] = IE[IU, IV] = -I[IU, IV] = [U, IV],$$

which shows $[U, IV] \in \mathfrak{g}_+$. Thus, $\mathfrak{g}_\pm$ are ideals in $\mathfrak{g}$ and the proposition follows from the following lemma. $\square$

Lemma 2.7. *Let $\mathfrak{g}$ admit a complex product structure $\{J, E\}$ and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be its associated Lie algebra. If both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are ideals in $\mathfrak{g}$ then $\mathfrak{g}$ is abelian.*

Proof. There is a splitting $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$, with $\mathfrak{g}_+$ and $\mathfrak{g}_- = J\mathfrak{g}_+$ ideals in $\mathfrak{g}$. Since $[\mathfrak{g}_+, \mathfrak{g}_-] = 0$, we have that $[JU, V] = [U, JV] = 0$ for all $U, V \in \mathfrak{g}_+$. From the integrability of J we obtain $J[U, V] = J[JU, JV]$, and then $[U, V] = [JU, JV] = 0$ for all $U, V \in \mathfrak{g}_+$. Hence $\mathfrak{g}_\pm$ are abelian ideals and it follows that $\mathfrak{g}$ is abelian. $\square$

Examples 2.8. (i) Given any Lie algebra $\mathfrak{u}$, set $\mathfrak{g} = \mathfrak{u} \times \mathfrak{u}$. Let $E : \mathfrak{g} \longrightarrow \mathfrak{g}$ be defined by $E(U, V) = (U, -V)$. Then E is clearly a product structure on $\mathfrak{g}$. However, Lemma 2.7 implies that there exists no complex product anticommuting with E unless $\mathfrak{u}$ is abelian. In this case any complex structure anticommuting with E can be seen to be equivalent to the standard complex structure on the abelian Lie algebra $\mathbb{C}^n = \mathbb{R}^n \times \mathbb{R}^n$ given by $J(U, V) = (-V, U)$.

(ii) The simplest non-abelian example of a Lie algebra carrying a complex product structure is given by $\mathfrak{aff}(\mathbb{R})$, the Lie algebra of the group $Aff(\mathbb{R})$ of affine motions of the real line. $\mathfrak{aff}(\mathbb{R})$ is a solvable non nilpotent Lie algebra with a basis $\{X, Y\}$ satisfying $[X, Y] = Y$. There is a complex structure J and a product structure E on $\mathfrak{aff}(\mathbb{R})$ given by $JX = Y$ and $EX = X$, $EY = -Y$. As $JE = -EJ$, $\{J, E\}$ defines a complex product structure on $\mathfrak{aff}(\mathbb{R})$.

An important class of Lie algebras carrying a complex product structure is obtained by the following construction. Consider a finite dimensional real vector space $\mathcal{A}$ equipped with a bilinear product $\mathcal{A} \times \mathcal{A} \longrightarrow \mathcal{A}$, $(a, b) \mapsto ab$. On the vector space $\mathcal{A} \oplus \mathcal{A}$ we consider the following skew-symmetric bilinear bracket:

$$(8) \quad [(a, b), (a', b')] = (aa' - a'a, ab' - a'b), \quad a, b, a', b' \in \mathcal{A}.$$

Suppose for a moment that the bracket (8) satisfies the Jacobi identity, hence defining a Lie algebra structure on $\mathcal{A} \oplus \mathcal{A}$. Let $J : \mathcal{A} \oplus \mathcal{A} \longrightarrow \mathcal{A} \oplus \mathcal{A}$ be the endomorphism given by

$$J(a, b) = (-b, a), \quad a, b \in \mathcal{A}.$$

A computation shows that J is a complex structure on $\mathcal{A} \oplus \mathcal{A}$. Furthermore, if we denote $\mathcal{A}_+ := \mathcal{A} \oplus \{0\}$ and $\mathcal{A}_- := \{0\} \oplus \mathcal{A}$, then it is easy to see that $\mathcal{A}_+$ is a subalgebra, $\mathcal{A}_-$ is an abelian ideal of $\mathcal{A} \oplus \mathcal{A}$ and also $\mathcal{A}_- = J\mathcal{A}_+$. Hence, letting $E : \mathcal{A} \oplus \mathcal{A} \rightarrow \mathcal{A} \oplus \mathcal{A}$ be given by $E|_{\mathcal{A}_+} = \mathbf{1}$, $E|_{\mathcal{A}_-} = -\mathbf{1}$, we obtain a complex product structure $\{J, E\}$ on $\mathcal{A} \oplus \mathcal{A}$.

Let us now investigate the conditions under which (8) does in fact satisfy the Jacobi identity. To do so, we shall need the following concept, which has already been intensively studied (see for instance [9, 10, 16, 29]).

Definition 2.9. A *left-symmetric algebra (LSA)* structure on a Lie algebra $\mathfrak{h}$ is a bilinear product $\mathfrak{h} \times \mathfrak{h} \rightarrow \mathfrak{h}$, $(x, y) \mapsto x \cdot y$, which satisfies

$$(9) \quad x \cdot (y \cdot z) - (x \cdot y) \cdot z = y \cdot (x \cdot z) - (y \cdot x) \cdot z$$

and

$$(10) \quad [x, y] = x \cdot y - y \cdot x.$$

Lemma 2.10. *The bracket on $\mathcal{A} \oplus \mathcal{A}$ given by (8) satisfies the Jacobi identity if and only if $\mathcal{A}$ is a Lie algebra with an LSA structure.*

Proof. Let $a, a', a'', b, b', b'' \in \mathcal{A}$. Then it is easy to see that

$$[[[a, b], (a', b')], (a'', b'')] + [[[a', b'], (a'', b'')], (a, b)] + [[[a'', b''], (a, b)], (a', b')] = 0$$

if and only if

$$(aa')a'' - a(a'a'') - (a'a)a'' + a'(aa'') + (a''a)a' - a''(aa') - (aa'')a' + a(a''a') + (a'a'')a - a'(a''a) - (a''a')a + a''(a'a) = 0$$

and

$$(11) \quad (aa')b'' - a(a'b'') - (a'a)b'' + a'(ab'') + (a''a)b' - a''(ab') - (aa'')b' + a(a''b') + (a'a'')b - a'(a''b) - (a''a')b + a''(a'b) = 0.$$

It is clear that if $\mathcal{A}$ is a Lie algebra with an LSA structure, then the Jacobi identity is valid on $\mathcal{A} \oplus \mathcal{A}$, using (9). Conversely, suppose that the bracket (8) satisfies the Jacobi identity. If we take $a' = b'' = 0$ and $a = x$, $a'' = y$, $b' = z$ in (11), we obtain equation (9). As we have already mentioned, $\mathcal{A}_+ = \mathcal{A} \oplus \{0\}$ is a Lie subalgebra of $\mathcal{A} \oplus \mathcal{A}$. Identifying $\mathcal{A}_+$ with $\mathcal{A}$ in the obvious way, $\mathcal{A}$ itself acquires a Lie algebra structure which satisfies $[a, a'] = aa' - a'a$. Hence $\mathcal{A}$ is a Lie algebra with an LSA structure and the proof of the lemma is complete. $\square$

Remarks. (i) LSA structures on a Lie algebra are also known as Koszul-Vinberg structures or affine structures.

(ii) If we define $(x, y, z) = x \cdot (y \cdot z) - (x \cdot y) \cdot z$ for all $x, y, z \in \mathfrak{h}$, condition (9) becomes $(x, y, z) = (y, x, z)$, whence the name “left-symmetric”.

(iii) Lemma 2.10 generalizes the result given in [6], where it is shown that the bracket (8) gives a Lie algebra structure on $\text{aff}(\mathcal{A}) := \mathcal{A} \oplus \mathcal{A}$ whenever $\mathcal{A}$ is an associative algebra. We will also use the notation $\text{aff}(\mathcal{A})$ for the Lie algebra $\mathcal{A} \oplus \mathcal{A}$ when $\mathcal{A}$ is a Lie algebra with an LSA structure.

To sum up, starting with a Lie algebra with an LSA structure, we have constructed a Lie algebra with a complex product structure in which one of the eigenspaces corresponding to the product structure is an (abelian) ideal. We now reveal the converse.

Proposition 2.11. *Let $(\mathfrak{g}, J, E)$ be a Lie algebra with a complex product structure, and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be its associated double Lie algebra, $\mathfrak{g}_- = J\mathfrak{g}_+$. Suppose $\mathfrak{g}_-$ is an ideal in $\mathfrak{g}$. Then $\mathfrak{g}_-$ is abelian and hence $\mathfrak{g}$ is isomorphic to the semidirect product $\mathfrak{g}_+ \ltimes_{\text{ad}_{\mathfrak{g}}} \mathfrak{g}_-$. Also, $\mathfrak{g}_+$ carries an LSA structure, which is given by*

$$X \cdot Y = -J[X, JY], \quad X, Y \in \mathfrak{g}_+$$

Proof. Let $X, Y \in \mathfrak{g}_-$. Since J is integrable, we have

$$(12) \quad J[X, Y] = [X, JY] + [JX, Y] + J[JX, JY].$$

Now, the left-hand side of (12) is in $\mathfrak{g}_+$, whereas the right-hand side is in $\mathfrak{g}_-$. So, both sides of (12) are zero, showing that $\mathfrak{g}_-$ is an *abelian* ideal. Then $\mathfrak{g}$ is isomorphic to the semidirect product $\mathfrak{g}_+ \ltimes_{\rho} \mathfrak{g}_-$, where the representation $\rho : \mathfrak{g}_+ \rightarrow \mathfrak{gl}(\mathfrak{g}_-)$ is simply given by $\rho(X) = \text{ad}(X)$, $X \in \mathfrak{g}_+$. To see that $\mathfrak{g}_+$ carries an LSA structure, we compute for $X, Y, Z \in \mathfrak{g}_+$:

$$\begin{aligned} X \cdot Y - Y \cdot X &= -J[X, JY] + J[Y, JX] \\ &= -J([X, JY] + [JX, Y]) \\ &= -J(J[X, Y] - J[JX, JY]) \\ &= [X, Y] \end{aligned}$$

using the integrability of J and the fact that $\mathfrak{g}_-$ is abelian. Thus, (10) holds. Also,

$$\begin{aligned} X \cdot (Y \cdot Z) - Y \cdot (X \cdot Z) &= -X \cdot (J[Y, JZ]) + Y \cdot (J[X, JZ]) \\ &= -J[X, [Y, JZ]] + J[Y, [X, JZ]] \\ &= J[JZ, [X, Y]] \quad (\text{using Jacobi}) \\ &= -J[[X, Y], JZ] \\ &= [X, Y] \cdot Z, \end{aligned}$$

from where (9) follows. $\square$

3. HYPERCOMPLEX STRUCTURES FROM COMPLEX PRODUCT STRUCTURES

In this section we will show that each complex product structure on a Lie algebra gives rise to a hypercomplex structure on the real Lie algebra underlying its complexification. Firstly, we recall

Definition 3.1. A *hypercomplex structure* on the Lie algebra $\mathfrak{g}$ is a pair $\{J_1, J_2\}$ of complex structures on $\mathfrak{g}$ satisfying $J_1 J_2 = -J_2 J_1$.

Note that $J_3 := J_1 J_2$ is another complex structure on $\mathfrak{g}$ and $\{J_1, J_2, J_3\}$ satisfy

$$J_1 J_2 = J_3, \quad J_2 J_3 = J_1, \quad J_3 J_1 = J_2.$$

Throughout this section, we shall use the notation: $\hat{\mathfrak{g}} = (\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$. We have already noted that multiplication by i in $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ defines a complex structure I on $\hat{\mathfrak{g}}$ satisfying (3). Now suppose that $\mathfrak{g}$ has dimension $2n$ and a complex structure J . In this situation, the

complexification of J can be regarded as a complex structure $\mathbb{J}$ on $\hat{\mathfrak{g}}$ that commutes with I . To see this, use the decomposition

$$\hat{\mathfrak{g}} = \mathfrak{g} \oplus I\mathfrak{g},$$

set $\mathbb{J}|_{\mathfrak{g}} = J$ and extend the definition of J to $I\mathfrak{g}$ by $\mathbb{J}(IX) = I(JX)$. It is easy to verify the integrability of $\mathbb{J}$, using (3).

Of course, I and $\mathbb{J}$ never determine a hypercomplex structure because they commute. Our aim is to construct hypercomplex structures on $\mathfrak{g}$ by retaining $\mathbb{J}$, but modifying I .

Lemma 3.2. *Let $\mathfrak{g}$ be a Lie algebra with a complex structure I satisfying (3). Suppose there is a splitting $\mathfrak{g} = \mathfrak{u}_1 \oplus \mathfrak{u}_2$ with $\mathfrak{u}_1, \mathfrak{u}_2$ complex subalgebras of $\mathfrak{g}$. Then the linear endomorphism $\mathbb{I}$ defined by*

$$\mathbb{I}|_{\mathfrak{u}_1} = I, \quad \mathbb{I}|_{\mathfrak{u}_2} = -I$$

is a complex structure on $\mathfrak{g}$.

Proof. It is clear that $\mathbb{I}^2 = -\mathbf{1}$, so, we only have to check the integrability of $\mathbb{I}$. Let $X = X_1 + X_2 \in \mathfrak{g}$, $Y = Y_1 + Y_2 \in \mathfrak{g}$, with $X_1, Y_1 \in \mathfrak{u}_1$ and $X_2, Y_2 \in \mathfrak{u}_2$. Then we have

$$\begin{aligned} \mathbb{I}[X, Y] &= \mathbb{I}[X_1 + X_2, Y_1 + Y_2] \\ &= I[X_1, Y_1] - I[X_2, Y_2] + \mathbb{I}([X_1, Y_2] + [X_2, Y_1]) \end{aligned}$$

On the other hand,

$$\begin{aligned} [\mathbb{I}X, Y] &= [IX_1 - IX_2, Y_1 + Y_2] \\ &= I[X_1, Y_1] + I[X_1, Y_2] - I[X_2, Y_1] - I[X_2, Y_2] \\ [X, \mathbb{I}Y] &= [X_1 + X_2, IY_1 - IY_2] \\ &= I[X_1, Y_1] - I[X_1, Y_2] + I[X_2, Y_1] - I[X_2, Y_2] \\ \mathbb{I}[\mathbb{I}X, \mathbb{I}Y] &= \mathbb{I}[IX_1 - IX_2, IY_1 - IY_2] \\ &= \mathbb{I}(-[X_1, Y_1] + [X_1, Y_2] + [X_2, Y_1] - [X_2, Y_2]) \\ &= -I[X_1, Y_1] + I[X_2, Y_2] + \mathbb{I}([X_1, Y_2] + [X_2, Y_1]) \end{aligned}$$

Hence, $\mathbb{I}[X, Y] = [\mathbb{I}X, Y] + [X, \mathbb{I}Y] + \mathbb{I}[\mathbb{I}X, \mathbb{I}Y]$ and $\mathbb{I}$ is integrable. $\square$

Theorem 3.3. *If $\mathfrak{g}$ has a complex product structure then $\hat{\mathfrak{g}}$ has a hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ with $\mathbb{I}$ and $\mathbb{J}$ as above.*

Proof. From Proposition 2.2, we have a decomposition $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$, with $\mathfrak{g}_+$ and $\mathfrak{g}_- = J\mathfrak{g}_+$ subalgebras of $\mathfrak{g}$. By complexifying, we obtain $\hat{\mathfrak{g}} = (\mathfrak{g}_+ \oplus I\mathfrak{g}_+) \oplus (\mathfrak{g}_- \oplus I\mathfrak{g}_-)$, where $\mathfrak{g}_+ \oplus I\mathfrak{g}_+$ and $\mathfrak{g}_- \oplus I\mathfrak{g}_-$ are complex subalgebras of $\hat{\mathfrak{g}}$. By the lemma, the endomorphism $\mathbb{I}$ defined by $\mathbb{I}|_{\mathfrak{g}_+ \oplus I\mathfrak{g}_+} = I$, $\mathbb{I}|_{\mathfrak{g}_- \oplus I\mathfrak{g}_-} = -I$ is a complex structure on $\hat{\mathfrak{g}}$. Also, it is clear that $\mathbb{I}$ and $\mathbb{J}$ anticommute. $\square$

Remark. It is easy to verify that two equivalent complex product structures on the Lie algebra $\mathfrak{g}$ give rise to equivalent hypercomplex structures on $\hat{\mathfrak{g}}$. We need only recall that two hypercomplex structures $\{J_k\}_{k=1,2}$ and $\{J'_k\}_{k=1,2}$ are said to be equivalent if there exists an automorphism ϕ of $\mathfrak{g}$ such that $\phi J_k = J'_k \phi$ for $k = 1, 2$.

Note that $\hat{\mathfrak{g}}$ also has a complex product structure, induced by the one on $\mathfrak{g}$. Indeed, define $\mathbb{E} : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}$ by $\mathbb{E} = -I\mathbb{I}$. As I commutes with $\mathbb{I}$, we have that $\mathbb{E}^2 = \mathbf{1}$ and $\mathbb{J}\mathbb{E} = -\mathbb{E}\mathbb{J}$. It is also easy to verify that $\mathbb{E}$ is integrable, as a consequence of (3). Finally, the eigenspaces corresponding to the eigenvalues $+1$ and -1 of $\mathbb{E}$ are, respectively, the subalgebras $\mathfrak{g}_+ \oplus I\mathfrak{g}_+$ and $\mathfrak{g}_- \oplus I\mathfrak{g}_-$. Hence, the pair $\{\mathbb{J}, \mathbb{E}\}$ determines a complex product structure on $\hat{\mathfrak{g}}$. Applying Theorem 3.3 to $\hat{\mathfrak{g}}$, we obtain another Lie algebra with a hypercomplex structure, with dimension twice the dimension of $\hat{\mathfrak{g}}$ and four times the dimension of $\mathfrak{g}$. In this way, starting with a $2n$ -dimensional Lie algebra $\mathfrak{g}$ endowed with a complex product structure, we obtain a family $\{\mathfrak{g}_{(k)} : k \in \mathbb{N}\}$ of Lie algebras carrying hypercomplex structures, where $\mathfrak{g}_{(1)} := \mathfrak{g}$, $\mathfrak{g}_{(k+1)} := \widehat{\mathfrak{g}_{(k)}}$ and then $\dim \mathfrak{g}_{(k)} = 2^{k+1}n$.

Examples 3.4. (i) If $\mathfrak{g} = \mathfrak{aff}(\mathbb{R})$, applying Theorem 3.3 we obtain a hypercomplex structure on $\hat{\mathfrak{g}} = \mathfrak{aff}(\mathbb{C})$, the Lie algebra of the group $Aff(\mathbb{C})$ of affine motions of the complex plane. $\mathfrak{aff}(\mathbb{C})$ has a basis $\{X, Y, \hat{X}, \hat{Y}\}$ with non-zero bracket relations

$$[X, Y] = -[\hat{X}, \hat{Y}] = Y, \quad [X, \hat{Y}] = [\hat{X}, Y] = \hat{Y},$$

and the hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ is given by

$$\begin{aligned} \mathbb{I}X &= \hat{X}, & \mathbb{I}Y &= -\hat{Y}, & \mathbb{I}^2 &= -\mathbf{1}, \\ \mathbb{J}X &= Y, & \mathbb{J}\hat{X} &= \hat{Y}, & \mathbb{J}^2 &= -\mathbf{1}. \end{aligned}$$

According to [4], $\mathfrak{aff}(\mathbb{C})$ is the only 4-dimensional Lie algebra with 2-dimensional commutator ideal admitting a hypercomplex structure.

(ii) More generally, recall from §2 that $\mathfrak{aff}(\mathcal{A})$ admits a complex product structure, where $\mathcal{A}$ is a Lie algebra with an LSA structure. Hence, Theorem 3.3 gives a hypercomplex structure on $\widehat{\mathfrak{aff}(\mathcal{A})} = \mathfrak{aff}(\mathcal{A}^{\mathbb{C}})$, where $\mathcal{A}^{\mathbb{C}}$ is the complex Lie algebra with an LSA structure obtained by complexifying $\mathcal{A}$. This hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ is given by

$$\mathbb{I}(a, b) = (ia, -ib), \quad \mathbb{J}(a, b) = (-b, a),$$

where $a, b \in \mathcal{A}^{\mathbb{C}}$. This hypercomplex structure coincides (up to a sign) with the hypercomplex structure constructed in [6] (where only associative algebras were considered).

We will prove next that the reductive Lie algebra $\mathfrak{gl}(2n, \mathbb{R})$ carries a canonical complex product structure; hence we will obtain a hypercomplex structure on $\mathfrak{gl}(2n, \mathbb{C})$.

Proposition 3.5. *$\mathfrak{gl}(2n, \mathbb{R})$ has a complex product structure for all $n \geq 1$.*

Proof. First let

$$(13) \quad E_0 = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix} \quad \text{and} \quad J_0 = \begin{pmatrix} 0 & -\mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}$$

denote the standard almost product and complex structures acting on $\mathbb{R}^{2n}$, where $\mathbf{1}$ is the $(n \times n)$ -identity matrix. Here E_0 and J_0 are expressed as $2n \times 2n$ matrices acting by left multiplication on column vectors. Then E_0 and J_0 define an almost product structure E and an almost complex structure J on the set $\mathfrak{gl}(2n, \mathbb{R})$ of $2n \times 2n$ matrices by right multiplication:

$$E(A) = AE_0, \quad J(A) = AJ_0$$

for any $A \in \mathfrak{gl}(2n, \mathbb{R})$. Clearly, $EJ = -JE$. To see that E is integrable, one can argue as follows.

Let $V = \mathbb{R}^{2n}$, so $\mathfrak{gl}(2n, \mathbb{R}) = \text{End}(V) = V \otimes V^*$. Multiplying two matrices corresponds to $(v \otimes \alpha) \cdot (w \otimes \beta) = \alpha(w)(v \otimes \beta)$ for $\alpha, \beta \in V^*$ and $v, w \in V$. On a generator $v \otimes \alpha \in V \otimes V^*$, the action of E is given by $E(v \otimes \alpha) = v \otimes E_0 \alpha$, where $E_0 \alpha = \alpha \circ E_0$. So, if $V_\pm^*$ denotes the (± 1) -eigenspace of E_0 on V^* , then the $(+1)$ - and (-1) -eigenspaces of E are given, respectively, by $V \otimes V_+^*$ and $V \otimes V_-^*$. It is easily seen that the product (and thus the Lie bracket) of two elements of $V \otimes V_+^*$ will produce a matrix that remains in that subspace, showing that $V \otimes V_+^*$ is a Lie subalgebra of $\text{End } V$. The same is true for $V \otimes V_-^*$ and hence E is integrable.

In the same way we may prove that J is integrable, by considering the $(\pm i)$ -eigenspaces of $J_0^\mathbb{C}$ on $V^\mathbb{C}$. Therefore, $\{J, E\}$ is a complex product structure on $\mathfrak{gl}(2n, \mathbb{R})$. $\square$

Corollary 3.6. *The induced hypercomplex structure on $\mathfrak{gl}(2n, \mathbb{C})$ is given by right multiplication by the matrices*

$$\mathbb{I} = \begin{pmatrix} i\mathbf{1} & 0 \\ 0 & -i\mathbf{1} \end{pmatrix}, \quad \mathbb{J} = J_0.$$

This hypercomplex structure on $\mathfrak{gl}(2n, \mathbb{C})$ merely corresponds to writing

$$(14) \quad \mathfrak{gl}(2n, \mathbb{C}) = (\mathbb{C}^{2n})^* \otimes \mathbb{H}^n \cong \bigoplus_{2n} \mathbb{H}^n,$$

and using right multiplication by unit quaternions on $\mathbb{H}^n$.

We close this section with a negative result, namely the failure of a naïve attempt to extend Proposition 3.5 to the Lie algebra

$$(15) \quad \mathfrak{sp}(n, \mathbb{R}) = \{X \in \mathfrak{gl}(2n, \mathbb{R}) : X^t J_0 + J_0 X = 0\}$$

of the symplectic group preserving a non-degenerate skew form on $\mathbb{R}^{2n}$ (notation of (13)).

Example 3.7. It can be seen that $X \in \mathfrak{sp}(n, \mathbb{R})$ if and only if

$$X = \begin{pmatrix} A & B \\ C & -A^t \end{pmatrix},$$

where $A, B, C \in \mathfrak{gl}(n, \mathbb{R})$ and B and C are symmetric. The subspace of $\mathfrak{sp}(n, \mathbb{R})$ given by

$$\left\{ \begin{pmatrix} A & 0 \\ 0 & -A^t \end{pmatrix} : A \in \mathfrak{gl}(n, \mathbb{R}) \right\}$$

is in fact a subalgebra of $\mathfrak{sp}(n, \mathbb{R})$ isomorphic to $\mathfrak{gl}(n, \mathbb{R})$. Also, the subspaces

$$\mathfrak{a}_+ = \left\{ \begin{pmatrix} 0 & B \\ 0 & 0 \end{pmatrix} : B \text{ symmetric} \right\}, \quad \mathfrak{a}_- = \left\{ \begin{pmatrix} 0 & 0 \\ C & 0 \end{pmatrix} : C \text{ symmetric} \right\}$$

are abelian subalgebras of $\mathfrak{sp}(n, \mathbb{R})$. We have thus obtained a decomposition

$$\mathfrak{sp}(n, \mathbb{R}) = \mathfrak{gl}(n, \mathbb{R}) \oplus \mathfrak{a}_+ \oplus \mathfrak{a}_-$$

of (15) into the direct sum of three subalgebras. The proof of the following result is based on simple matrix computations best left to the reader.

Proposition 3.8. *There is no complex product structure $\{J, E\}$ on $\mathfrak{sp}(n, \mathbb{R})$ such that $J\mathfrak{a}_+ = \mathfrak{a}_-$.*

4. COMPLEX PRODUCT STRUCTURES FROM MATCHED PAIRS OF LIE ALGEBRAS

In this section we will characterize Lie algebras carrying a complex product structure in terms of double Lie algebras and matched pairs of Lie algebras endowed with a left-symmetric algebra (LSA) structure.

In Proposition 2.5, we characterized those double Lie algebras associated to a complex product structure. Given a double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$, the existence of such a structure on $\mathfrak{g}$ depends on the existence of a linear isomorphism $\varphi : \mathfrak{g}_+ \longrightarrow \mathfrak{g}_-$ which satisfies a certain property that involves the Lie bracket of $\mathfrak{g}$, not only the brackets of $\mathfrak{g}_+$ and $\mathfrak{g}_-$. On the other hand, we should like to construct Lie algebras with a complex product structure beginning with two Lie algebras of the same dimension, which are not a priori subalgebras of another one. The idea is to obtain certain conditions on these two Lie algebras which ensure that their direct sum as vector spaces admits a Lie bracket with respect to which both summands are Lie subalgebras.

Firstly, let us recall the following construction, which appears for example in [22, 23]. Let $(\mathfrak{u}, \mathfrak{v}, \rho, \mu)$ be a *matched pair of Lie algebras*, i.e. $\mathfrak{u}$ and $\mathfrak{v}$ are Lie algebras and $\rho : \mathfrak{u} \longrightarrow \mathfrak{gl}(\mathfrak{v})$ and $\mu : \mathfrak{v} \longrightarrow \mathfrak{gl}(\mathfrak{u})$ are representations satisfying

$$(16) \quad \rho(X)[A, B] - [\rho(X)A, B] - [A, \rho(X)B] + \rho(\mu(A)X)B - \rho(\mu(B)X)A = 0,$$

$$(17) \quad \mu(A)[X, Y] - [\mu(A)X, Y] - [X, \mu(A)Y] + \mu(\rho(X)A)Y - \mu(\rho(Y)A)X = 0,$$

for $X, Y \in \mathfrak{u}$ and $A, B \in \mathfrak{v}$. In this case, the vector space $\mathfrak{g} = \mathfrak{u} \oplus \mathfrak{v}$ admits a Lie bracket, given by

$$[(X, A), (Y, B)] = ([X, Y] + \mu(A)Y - \mu(B)X, [A, B] + \rho(X)B - \rho(Y)A),$$

for $X, Y \in \mathfrak{u}$ and $A, B \in \mathfrak{v}$. We will denote this new Lie algebra by $\mathfrak{g} = \mathfrak{u} \bowtie_{\mu}^{\rho} \mathfrak{v}$ (or simply $\mathfrak{g} = \mathfrak{u} \bowtie \mathfrak{v}$) and will call it the *bicrossproduct* of $\mathfrak{u}$ and $\mathfrak{v}$. Observe that $\mathfrak{u} \equiv \mathfrak{u} \oplus \{0\}$ and $\mathfrak{v} \equiv \{0\} \oplus \mathfrak{v}$ are subalgebras of $\mathfrak{g}$ and then $(\mathfrak{u} \bowtie \mathfrak{v}, \mathfrak{u}, \mathfrak{v})$ is a double Lie algebra.

Conversely, if $\mathfrak{u}$ and $\mathfrak{v}$ are Lie subalgebras of a Lie algebra $\mathfrak{g}$ such that $\mathfrak{u} \oplus \mathfrak{v} = \mathfrak{g}$, i.e. $(\mathfrak{g}, \mathfrak{u}, \mathfrak{v})$ is a double Lie algebra, then $(\mathfrak{u}, \mathfrak{v}, \rho, \mu)$ forms a matched pair of Lie algebras, where the representations $\rho : \mathfrak{u} \longrightarrow \mathfrak{gl}(\mathfrak{v})$ and $\mu : \mathfrak{v} \longrightarrow \mathfrak{gl}(\mathfrak{u})$ are determined by

$$(18) \quad [X, A] = -\mu(A)X + \rho(X)A, \quad X \in \mathfrak{u}, A \in \mathfrak{v},$$

so that $\mathfrak{g} = \mathfrak{u} \bowtie \mathfrak{v}$.

Let us now suppose that $\mathfrak{g}$ is a Lie algebra endowed with a complex product structure $\{J, E\}$ and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be its associated double Lie algebra, where $\mathfrak{g}_- = J\mathfrak{g}_+$. From the paragraph above, there exist representations $\rho : \mathfrak{g}_+ \longrightarrow \mathfrak{gl}(\mathfrak{g}_-)$ and $\mu : \mathfrak{g}_- \longrightarrow \mathfrak{gl}(\mathfrak{g}_+)$ such that $(\mathfrak{g}_+, \mathfrak{g}_-, \rho, \mu)$ forms a matched pair of Lie algebras and then $\mathfrak{g} = \mathfrak{g}_+ \bowtie \mathfrak{g}_-$. In this situation, a particular feature is given by the existence of the isomorphism $J : \mathfrak{g}_+ \longrightarrow \mathfrak{g}_-$, since it allows us to produce a pair of representations equivalent to ρ and μ in the following way. Let $\tilde{\rho} : \mathfrak{g}_+ \longrightarrow \mathfrak{gl}(\mathfrak{g}_+)$ and $\tilde{\mu} : \mathfrak{g}_- \longrightarrow \mathfrak{gl}(\mathfrak{g}_-)$ be given by

$$(19) \quad \tilde{\rho}(X) := -J\rho(X)J, \quad \tilde{\mu}(A) := -J\mu(A)J,$$

for $X \in \mathfrak{g}_+$ and $A \in \mathfrak{g}_-$. It is a simple matter to check that $\tilde{\rho}$ and $\tilde{\mu}$ are indeed representations of $\mathfrak{g}_+$ and $\mathfrak{g}_-$, respectively. With this notation, (18) translates into

$$[X, A] = -J\tilde{\rho}(X)JA + J\tilde{\mu}(A)JX,$$

for $X \in \mathfrak{g}_+$ and $A \in \mathfrak{g}_-$. So, if $\pi_+ : \mathfrak{g} \longrightarrow \mathfrak{g}_+$ and $\pi_- : \mathfrak{g} \longrightarrow \mathfrak{g}_-$ denote the projections, we have the relations

$$\tilde{\rho}(X)Y = -\pi_+ J[X, JY], \quad \tilde{\mu}(A)B = -\pi_- J[A, JB],$$

for $X, Y \in \mathfrak{g}_+$ and $A, B \in \mathfrak{g}_-$.

The relationship between Lie algebras with a complex product structure and Lie algebras with an LSA is the content of the following result, which is a generalization of Proposition 2.11.

Theorem 4.1. *Let $(\mathfrak{g}, J, E)$ be a Lie algebra with a complex product structure and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be its associated double Lie algebra with $\mathfrak{g}_- = J\mathfrak{g}_+$. Then $\mathfrak{g}_+$ and $\mathfrak{g}_-$ carry an LSA structure.*

Proof. Define a bilinear product on $\mathfrak{g}_+$ in the following manner

$$\mathfrak{g}_+ \times \mathfrak{g}_+ \longrightarrow \mathfrak{g}_+, \quad (X, Y) \mapsto X \cdot Y := \tilde{\rho}(X)Y,$$

where $\tilde{\rho}$ is as in (19). Let us first verify (10). For $X, Y \in \mathfrak{g}_+$, we have

$$\begin{aligned} X \cdot Y - Y \cdot X &= \tilde{\rho}(X)Y - \tilde{\rho}(Y)X \\ &= -\pi_+ J([X, JY] + [JX, Y]) \\ &= -\pi_+ J(J[X, Y] - J[JX, JY]) \\ &= \pi_+ ([X, Y] - [JX, JY]) \\ &= [X, Y]. \end{aligned}$$

The third equality follows from the integrability of J . Next, to verify (9), we perform the following computation. For $X, Y, Z \in \mathfrak{g}_+$,

$$X \cdot (Y \cdot Z) - Y \cdot (X \cdot Z) = \tilde{\rho}(X)\tilde{\rho}(Y)Z - \tilde{\rho}(Y)\tilde{\rho}(X)Z = \tilde{\rho}([X, Y])Z = [X, Y] \cdot Z$$

since $\tilde{\rho}$ is a representation. Also,

$$(X \cdot Y) \cdot Z - (Y \cdot X) \cdot Z = (X \cdot Y - Y \cdot X)Z = [X, Y] \cdot Z.$$

Hence, $X \cdot (Y \cdot Z) - Y \cdot (X \cdot Z) = (X \cdot Y) \cdot Z - (Y \cdot X) \cdot Z$ and this implies (9).

To show that $\mathfrak{g}_-$ carries an LSA structure too, consider the bilinear product

$$\mathfrak{g}_- \times \mathfrak{g}_- \longrightarrow \mathfrak{g}_-, \quad (A, B) \mapsto A \cdot B := \tilde{\mu}(A)B,$$

where $\tilde{\mu}$ is as in (19). In the same way as before, it is shown that this product satisfies (9) and (10). $\square$

Remarks. (i) A semisimple Lie algebra does not admit any LSA structure (see [10, 16]). Thus $\mathfrak{g}_+$ and $\mathfrak{g}_-$ cannot be semisimple.

(ii) There do exist compact Lie algebras with an LSA structure. The simplest is $\mathfrak{so}(3) \oplus \mathbb{R}$, the Lie algebra of the Lie groups $U(2)$ and $S^3 \times S^1$. An LSA structure on $\mathfrak{so}(3) \oplus \mathbb{R}$ is obtained from the associative algebra $\mathbb{H}$ of quaternions. (This is related to the fact that $S^3 \times S^1$ is an affine quotient of $\mathbb{H} \setminus \{0\}$, and admits a left-invariant flat torsion-free connection [5]; see §5).

(ii) Many, but not all, nilpotent Lie algebras admit LSA structures [7].

Conversely, we shall show that Lie algebras arising from certain pairs of Lie algebras with LSA structures, admit a complex product structure.

Theorem 4.2. *Let $\mathfrak{u}$ and $\mathfrak{v}$ be two Lie algebras of the same dimension, both carrying an LSA structure. Suppose there exists a linear isomorphism $\varphi : \mathfrak{u} \longrightarrow \mathfrak{v}$ such that the representations $\rho : \mathfrak{u} \longrightarrow \mathfrak{gl}(\mathfrak{v})$ and $\mu : \mathfrak{v} \longrightarrow \mathfrak{gl}(\mathfrak{u})$ given by*

$$\rho(X)A = \varphi(X \cdot \varphi^{-1}(A)), \quad \mu(A)X = \varphi^{-1}(A \cdot \varphi(X))$$

satisfy (16) and (17). Then $(\mathfrak{u}, \mathfrak{v}, \rho, \mu)$ forms a matched product of Lie algebras and the bicrossproduct Lie algebra $\mathfrak{g} = \mathfrak{u} \bowtie \mathfrak{v}$ admits a complex product structure.

Proof. We start by defining an endomorphism $E : \mathfrak{g} \longrightarrow \mathfrak{g}$ by $E|_{\mathfrak{u}} = \mathbf{1}$, $E|_{\mathfrak{v}} = -\mathbf{1}$. Then $E^2 = \mathbf{1}$ and since the corresponding eigenspaces are Lie subalgebras of $\mathfrak{g}$, E determines a product structure on $\mathfrak{g}$. Next, we define another endomorphism J of $\mathfrak{g}$ by

$$J(X, A) = (-\varphi^{-1}(A), \varphi(X)), \quad X \in \mathfrak{u}, A \in \mathfrak{v}.$$

It is clear that $J^2 = -\mathbf{1}$. We only have to prove the integrability. In order to do so, we will consider several cases.

(i) For $X, Y \in \mathfrak{u}$, we have:

$$J[(X, 0), (Y, 0)] = J([X, Y], 0) = (0, \varphi[X, Y]).$$

On the other hand,

$$\begin{aligned} [J(X, 0), (Y, 0)] &= [(0, \varphi X), (Y, 0)] \\ &= -(-\mu(\varphi X)Y, \rho(Y)\varphi X) \\ &= (\varphi^{-1}(\varphi X \cdot \varphi Y), -\varphi(Y \cdot X)), \\ [(X, 0), J(Y, 0)] &= [(X, 0), (0, \varphi Y)] \\ &= (-\mu(\varphi Y)X, \rho(X)\varphi Y) \\ &= (-\varphi^{-1}(\varphi Y \cdot \varphi X), \varphi(X \cdot Y)) \end{aligned}$$

and

$$\begin{aligned} J[J(X, 0), J(Y, 0)] &= J[(0, \varphi X), (0, \varphi Y)] \\ &= J(0, [\varphi X, \varphi Y]) \\ &= (-\varphi^{-1}[\varphi X, \varphi Y], 0). \end{aligned}$$

Therefore, using (10), we obtain

$$J[(X, 0), (Y, 0)] = [J(X, 0), (Y, 0)] + [(X, 0), J(Y, 0)] + J[J(X, 0), J(Y, 0)].$$

(ii) For $A, B \in \mathfrak{v}$, with a similar computation to the one above, we obtain

$$J[(0, A), (0, B)] = [J(0, A), (0, B)] + [(0, A), J(0, B)] + J[J(0, A), J(0, B)].$$

(iii) Finally, for $X \in \mathfrak{u}$, $A \in \mathfrak{v}$, one obtains

$$\begin{aligned} J[(X, 0), (0, A)] &= J(-\mu(A)X, \rho(X)A) \\ &= (-\varphi^{-1}(\rho(X)A), -\varphi(\mu(A)X)) \\ &= (-X \cdot \varphi^{-1}A, -A \cdot \varphi X), \\ [J(X, 0), (0, A)] &= [(0, \varphi X), (0, A)] = (0, [\varphi X, A]), \\ [(X, 0), J(0, A)] &= [(X, 0), (-\varphi^{-1}A, 0)] = (-[X, \varphi^{-1}A], 0), \end{aligned}$$

and also

$$\begin{aligned}
J[J(X, 0), J(0, A)] &= J[(0, \varphi X), (-\varphi^{-1}A, 0)] \\
&= -J[(-\varphi^{-1}A, 0), (0, \varphi X)] \\
&= -J(\mu(\varphi X)\varphi^{-1}A, -\rho(\varphi^{-1}A)\varphi X) \\
&= -J(\varphi^{-1}(\varphi X \cdot A), -\varphi(\varphi^{-1}A \cdot X)) \\
&= (-\varphi^{-1}A \cdot X, -\varphi X \cdot A).
\end{aligned}$$

Using (10) again, we obtain

$$J[(X, 0), (0, A)] = [J(X, 0), (0, A)] + [(X, 0), J(0, A)] + J[J(X, 0), J(0, A)].$$

This concludes the verification of the integrability of J . Thus, J is a complex structure which anticommutes with E , since $\mathfrak{v} = J\mathfrak{u}$, showing that $\{J, E\}$ is a complex product structure on $\mathfrak{g}$. $\square$

Remarks. (i) Suppose that the LSA structure on $\mathfrak{v}$ is trivial, i.e. $A \cdot B = 0$ for all $A, B \in \mathfrak{v}$. Hence $\mathfrak{v}$ is an abelian ideal of $\mathfrak{u} \ltimes \mathfrak{v}$ and, in fact, $\mathfrak{u} \ltimes \mathfrak{v} \cong \mathfrak{aff}(\mathfrak{u})$.

(ii) Majid [22] has constructed for the Lie algebra $\mathfrak{g}$ of a compact semisimple Lie group another Lie algebra $\mathfrak{g}'$ such that $(\mathfrak{g}, \mathfrak{g}')$ determines a matched pair of Lie algebras. We have shown that for each Lie algebra carrying an LSA structure there exists another Lie algebra of the same dimension (namely, the abelian one) such that they determine a matched pair of Lie algebras.

Corollary 4.3. *Consider the abelian Lie algebra $\mathbb{R}^n$ endowed with an LSA structure. Then $(\mathbb{R}^n, \mathbb{R}^n, \rho, \rho)$ is a matched pair of Lie algebras, where $\rho : \mathbb{R}^n \longrightarrow \mathfrak{gl}(\mathbb{R}^n)$ is the representation of $\mathbb{R}^n$ given by $\rho(X)Y = X \cdot Y$. Hence, $\mathfrak{g} = \mathbb{R}^n \ltimes \mathbb{R}^n$ admits a complex product structure.*

Proof. Simply set $\varphi = \mathbf{1}_{\mathbb{R}^n}$ in Theorem 4.2. It is trivial to verify that (16) and (17) hold and then the corollary follows. $\square$

The previous corollary can be generalized in the following way.

Corollary 4.4. *Suppose the abelian Lie algebra $\mathbb{R}^n$ carries two LSA structures, denoted by $\mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}^n, (X, Y) \mapsto X \cdot Y$ and $\mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}^n, (X, Y) \mapsto X * Y$. Denote by $L : \mathbb{R}^n \longrightarrow \mathfrak{gl}(\mathbb{R}^n)$ and $L^* : \mathbb{R}^n \longrightarrow \mathfrak{gl}(\mathbb{R}^n)$ the representations given by left multiplications: $L_X Y = X \cdot Y, \quad L_X^* Y = X * Y$. If*

$$L_X L_Y^* = L_Y L_X^*, \quad L_X^* L_Y = L_Y^* L_X$$

for all $X, Y \in \mathbb{R}^n$, then $(\mathbb{R}^n, \mathbb{R}^n, L, L^)$ is a matched pair of Lie algebras and hence, $\mathfrak{g} = \mathbb{R}^n \ltimes \mathbb{R}^n$ carries a complex product structure.*

Proof. Take $\varphi = \mathbf{1}_{\mathbb{R}^n}$ in Theorem 4.2. $\square$

Let $(\mathfrak{g}, J, E)$ be a Lie algebra with a complex product structure and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be its associated double Lie algebra. We have shown that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ carry LSA structures. A natural question to ask is whether $\mathfrak{g}$ itself supports an LSA structure, which restricts

to the ones already existing on $\mathfrak{g}_\pm$. We can naturally extend the bilinear products on $\mathfrak{g}_+$ and $\mathfrak{g}_-$ to $\mathfrak{g}$ by using the representations $\rho, \mu, \tilde{\rho}, \tilde{\mu}$ in the following fashion:

$$(X + A) \cdot (Y + B) = \tilde{\rho}(X)Y + \rho(X)B + \mu(A)Y + \tilde{\mu}(A)B, \quad X, Y \in \mathfrak{g}_+, A, B \in \mathfrak{g}_-,$$

or more simply

$$(20) \quad (X + A) \cdot (Y + B) = X \cdot Y + \rho(X)B + \mu(A)Y + A \cdot B$$

where we are using the LSA structure on $\mathfrak{g}_\pm$. It is a simple matter to verify that this product on $\mathfrak{g}$ verifies (10). However, it does not necessarily fulfill (9). We will give an example of this fact in the next section (see Example 6.3).

To see when (20) satisfies (9), we introduce the following notation. Define $\Phi : \mathfrak{g}_+ \longrightarrow (\mathfrak{g}_- \otimes \mathfrak{g}_-)^* \otimes \mathfrak{g}_-$ by

$$(21) \quad \Phi(X)(A, B) = \rho(X)(A \cdot B) - (\rho(X)A) \cdot B - A \cdot (\rho(X)B) + \rho(\mu(A)X)B$$

and $\Psi : \mathfrak{g}_- \longrightarrow (\mathfrak{g}_+ \otimes \mathfrak{g}_+)^* \otimes \mathfrak{g}_+$ by

$$(22) \quad \Psi(A)(X, Y) = \mu(A)(X \cdot Y) - (\mu(A)X) \cdot Y - X \cdot (\mu(A)Y) + \mu(\rho(X)A)Y,$$

for $X, Y \in \mathfrak{g}_+, A, B \in \mathfrak{g}_-$. Then equations (16) and (17) can be rewritten simply as

$$\Phi(X)(A, B) = \Phi(X)(B, A), \quad \Psi(A)(X, Y) = \Psi(A)(Y, X),$$

for $X, Y \in \mathfrak{g}_+, A, B \in \mathfrak{g}_-$. Using Lemma 3.0.3 of [11], we obtain that the product defined above determines an LSA structure on $\mathfrak{g}$ if and only if $\Phi \equiv 0$ and $\Psi \equiv 0$.

Summing up,

Proposition 4.5. *Let $(\mathfrak{g}, J, E)$ be a Lie algebra with a complex product structure with associated double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ and $\mathfrak{g}_- = J\mathfrak{g}_+$. There exists an LSA structure on $\mathfrak{g}$ extending the LSA structures on $\mathfrak{g}_+$ and $\mathfrak{g}_-$ if and only if $\Phi \equiv 0$ and $\Psi \equiv 0$, where Φ and Ψ are as in (21) and (22).*

Corollary 4.6. *Let $(\mathfrak{g}, J, E)$ be a Lie algebra with a complex product structure and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be its associated double Lie algebra, with $\mathfrak{g}_- = J\mathfrak{g}_+$. Suppose that $\mathfrak{g}_-$ is an ideal of $\mathfrak{g}$. Then $\mathfrak{g}$ admits an LSA structure which extends the LSA structures on $\mathfrak{g}_+$ and $\mathfrak{g}_-$.*

Proof. We already know that in this case the LSA structure on $\mathfrak{g}_-$ is trivial (i.e. $A \cdot B = 0$ for all $A, B \in \mathfrak{g}_-$) and consequently, the μ is also trivial. Substituting into (21) and (22), we get $\Phi \equiv 0$ and $\Psi \equiv 0$. Hence $\mathfrak{g}$ does admit an LSA structure. $\square$

5. TORSION-FREE CONNECTIONS AND G -STRUCTURES

We start recalling that for an arbitrary connection ∇ on (the tangent bundle of) a manifold M , the torsion and curvature tensor fields T and R are defined by

$$T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]$$

$$R(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]}$$

for X, Y smooth vector fields on M . The connection is called *torsion-free* when $T = 0$, and *flat* when $R = 0$. A flat torsion-free affine connection on M gives rise to an *affine structure* on M [2, 24].

Let G be a Lie group with Lie algebra $\mathfrak{g}$ and suppose that G admits a left-invariant connection ∇ . This means that if $X, Y \in \mathfrak{g}$ are two left-invariant vector fields on G then $\nabla_X Y \in \mathfrak{g}$ is also left-invariant. Accordingly, one may define a connection on a Lie algebra $\mathfrak{g}$ to be merely a $\mathfrak{g}$ -valued bilinear form $\mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g}$. One can speak of the torsion and curvature of such a connection using the formulae above, with brackets determined by the structure of $\mathfrak{g}$. We can now re-interpret the results obtained in previous sections in terms of connections.

Let $\mathfrak{g}$ be a Lie algebra with a complex product structure $\{J, E\}$ and let $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ be the associated double Lie algebra. In §4, we have shown that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ carry an LSA structure, and we introduced a bilinear product on $\mathfrak{g}$ which extends these LSA structures and satisfies (10). This product becomes a torsion-free affine connection ∇^{CP} (CP for ‘complex product’) on $\mathfrak{g}$ by setting

$$\nabla_X^{\text{CP}} Y = X \cdot Y,$$

for $X, Y \in \mathfrak{g}$. The torsion-free condition reads

$$X \cdot Y - Y \cdot X = [X, Y],$$

and the connection is flat if

$$X \cdot (Y \cdot Z) - Y \cdot (X \cdot Z) = [X, Y] \cdot Z.$$

Since this is exactly condition (9), it follows that ∇^{CP} restricts to *flat* torsion-free connections on $\mathfrak{g}_+$ and $\mathfrak{g}_-$. In general though, ∇^{CP} is not itself flat (see Example 6.3 in §6).

Proposition 5.1. *If $(\mathfrak{g}, J, E)$ is a Lie algebra carrying a complex product structure then $\nabla^{\text{CP}} J = 0 = \nabla^{\text{CP}} E$. Moreover, ∇^{CP} is the unique torsion-free connection on $\mathfrak{g}$ for which J and E are parallel.*

Proof. The parallelism of J, E is quickly established from the definitions of the representations $\rho, \mu, \tilde{\rho}, \tilde{\mu}$ and equation (20).

As to uniqueness, suppose ∇, ∇' are two connections on $\mathfrak{g}$ which satisfy the conditions in the statement. Define $A : \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g}$ by $A(X, Y) = \nabla_X Y - \nabla'_X Y$. Since ∇, ∇' are both torsion-free,

$$\begin{aligned} A(Y, X) &= \nabla_Y X - \nabla'_Y X \\ &= \nabla_X Y + [Y, X] - \nabla'_X Y + [X, Y] = A(X, Y), \end{aligned}$$

and A is symmetric. Since $\nabla J = 0$, we obtain

$$\begin{aligned} A(X, JY) &= \nabla_X JY - \nabla'_X JY \\ &= J(\nabla_X Y - \nabla'_X Y) = JA(X, Y) \end{aligned}$$

and (by symmetry) $A(JX, Y) = JA(X, Y)$. Since also $\nabla E = 0$, the following is valid:

$$A(X, EY) = EA(X, Y) = A(EX, Y).$$

Then, $A(JX, EY) = EA(JX, Y) = JEA(X, Y)$, and

$$A(JX, EY) = JA(X, EY) = EJA(X, Y) = -JEA(X, Y).$$

It follows that $A(X, Y) = 0$ for all $X, Y \in \mathfrak{g}$, and so $\nabla = \nabla'$. □

We have shown in Theorem 3.3 that the complex product structure $\{J, E\}$ on $\mathfrak{g}$ determines a hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ on $\hat{\mathfrak{g}} = (\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$. In this case, any Lie group G with $L(G) = \hat{\mathfrak{g}}$ is a *hypercomplex manifold*, since it admits a pair $\{\mathbb{I}, \mathbb{J}\}$ of anticommuting complex structures. These are obtained by left-translating those that are already defined on $\mathfrak{g} \cong T_e G$. Every hypercomplex structure on M uniquely determines a torsion-free connection ∇^{HC} (HC for ‘hypercomplex’), sometimes called the *Obata connection*, with respect to which $\mathbb{I}, \mathbb{J}$ are parallel. Thus,

$$\nabla^{\text{HC}} \mathbb{I} = 0, \quad \nabla^{\text{HC}} \mathbb{J} = 0$$

(see [25]). Since $\mathbb{I}, \mathbb{J}$ are left-invariant, it follows that ∇^{HC} is left-invariant. In this way, we obtain a torsion-free connection on $\hat{\mathfrak{g}}$, that we also call the Obata connection.

On the other hand, the connection ∇^{CP} on $\mathfrak{g}$ extends naturally to a torsion-free connection $\hat{\nabla}^{\text{CP}}$ on $\hat{\mathfrak{g}}$ by setting

$$\hat{\nabla}_{X+IY}^{\text{CP}}(X' + IY') = (\nabla_X^{\text{CP}} X' - \nabla_Y^{\text{CP}} Y') + I(\nabla_X^{\text{CP}} Y' + \nabla_Y^{\text{CP}} X'),$$

for $X, X', Y, Y' \in \mathfrak{g}$. Since $\nabla^{\text{CP}} J = 0$, it is clear that $\hat{\nabla}^{\text{CP}} \mathbb{J} = 0$, and a simple computation shows that $\hat{\nabla}^{\text{CP}} \mathbb{I} = 0$. By uniqueness of the Obata connection, we have $\hat{\nabla}^{\text{CP}} = \nabla^{\text{HC}}$. We have thus proven

Proposition 5.2. *The canonical extension of ∇^{CP} to $\hat{\mathfrak{g}}$ coincides with the connection ∇^{HC} determined by the hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ on $\hat{\mathfrak{g}}$ given by Theorem 3.3.*

Corollary 5.3. *The Obata connection ∇^{HC} on $\hat{\mathfrak{g}}$ is flat if and only if the complex product connection ∇^{CP} on $\mathfrak{g}$ is flat.*

Example 5.4. The corollary applies to $\mathfrak{g} = \mathfrak{gl}(2n, \mathbb{R})$. For the proof of Proposition 3.5 shows that the complex product structure on $\mathfrak{gl}(2n, \mathbb{R})$ realizes it as $2n$ copies of the flat structure on $\mathbb{R}^{2n}$. In the same way, the induced hypercomplex structure on $\mathfrak{gl}(2n, \mathbb{C})$ is $2n$ copies (14) of flat space $\mathbb{H}^n$.

Recall the matrices J_0, E_0 defined in (13), and acting by left multiplication on the space $\mathbb{R}^{2n}$ of column vectors. The subgroup of $GL(2n, \mathbb{R})$ consisting of matrices commuting with J can be identified with $GL(n, \mathbb{C})$. The subgroup of those commuting with both J and E consists of invertible block-diagonal matrices

$$(23) \quad \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}$$

and can be identified with $GL(n, \mathbb{R})$.

We can generalize the preceding theory by

Definition 5.5. A *complex product manifold* is a smooth manifold M , of even dimension $2n$, equipped with a $GL(n, \mathbb{R})$ -structure (relative to the above inclusion $GL(n, \mathbb{R}) \subset GL(2n, \mathbb{R})$) admitting a $GL(n, \mathbb{R})$ connection.

In the general context of G -structures, there is a well-known series of obstructions to a given structure being equivalent to the standard structure on flat space [31]. The existence of a torsion-free connection is in general only the first such obstruction. The proof of Proposition 5.1 can be extended to show that a torsion-free $GL(n, \mathbb{R})$ -connection will (if it exists) be unique. Full details of this and the subsequent remarks appear in [1].

Given a $GL(n, \mathbb{R})$ -structure, we can regard its Lie algebra $\mathfrak{gl}(n, \mathbb{R})$ as a subspace of $\text{End } T = T \otimes T^*$, where $T = \mathbb{R}^{2n}$ stands for the tangent space representation. Torsion properties are determined by the natural mapping

$$f_1 : \mathfrak{gl}(n, \mathbb{R}) \otimes T^* \rightarrow T \otimes \bigwedge^2 T^*.$$

The difference of two compatible torsion-free connections is a tensor that hypothetically takes values at each point in $\ker f_1$, which is the so-called the first prolongation of $\mathfrak{gl}(n, \mathbb{R})$ (as a subalgebra of $\mathfrak{gl}(2n, \mathbb{R})$). The above remarks then imply

Lemma 5.6. $\ker f_1 = \{0\}$.

Let $T = T_+ \oplus T_-$ be the decomposition reflecting (23), so that $T_+ \cong T_-$ is the standard representation of $GL(n, \mathbb{R})$. The torsion-free condition ensures that the distributions determined by T_+ , T_- are integrable and totally geodesic. Moreover, M has an integrable complex structure J for which $T_- = JT_+$. The spaces $T_{\pm}$ now play the role of $\mathfrak{g}_{\pm}$ in §4.

The second (and only remaining) obstruction to flatness of the $GL(n, \mathbb{R})$ -structure of M is provided by the curvature tensor R of ∇ . This takes values in the kernel of the linear mapping

$$f_2 : \mathfrak{gl}(n, \mathbb{R}) \otimes \bigwedge^2 T^* \rightarrow T \otimes \bigwedge^3 T^*$$

that establishes the first Bianchi identity. The existence of an LSA structure on $\mathfrak{g}_{\pm}$ can be seen as a special case of the next result, that can be proved by working in terms of the standard $GL(n, \mathbb{R})$ -module $V = \mathbb{R}^n$, using the isomorphisms $T_+ \cong V \cong T_-$ and $\mathfrak{gl}(n, \mathbb{R}) \cong V^* \otimes V$ [1].

Lemma 5.7. *The restriction of R to $\mathfrak{gl}(n, \mathbb{R}) \otimes \bigwedge^2 T_{\pm}^*$ vanishes.*

This implies that the restriction of ∇ to each of the integrable submanifolds tangent to $T_{\pm}$ is flat.

The symmetries of R resemble those of the curvature tensor of a hypercomplex manifold (described in [27]), which is not surprising given the intimate relationship between the two structures. Hypercomplex structures are associated to the group $GL(n, \mathbb{H})$ in the same way that complex product structures are associated to $GL(n, \mathbb{R})$. The two groups can be extended to $GL(n, \mathbb{R}) \times GL(2, \mathbb{R})$ and $GL(n, \mathbb{H}) \times GL(1, \mathbb{H})$ respectively, and the resulting G -structures are both types of paraconformal geometries [3].

The above manifold theory can of course be applied to any Lie group G whose Lie algebra $\mathfrak{g}$ admits a complex product structure $\{J, E\}$. We may regard J, E as tensors on G by left-translating the ones on $\mathfrak{g} = T_e G$. If $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ is the double Lie algebra defined by E , let G_+ and G_- be Lie subgroups of G such that $L(G_{\pm}) = \mathfrak{g}_{\pm}$. We learn from [21] that (G, G_+, G_-) is a local double Lie group, i.e. the mapping $G_+ \times G_- \rightarrow G$ defined by $(g_+, g_-) \mapsto g_+ g_-$ is a local diffeomorphism near the identities. Since $\mathfrak{g}_{\pm}$ carry LSA structures, both G_+ and G_- are flat affine Lie groups, whilst G itself need not be flat.

In the notation of §2, let $\hat{G}$ be a Lie group with Lie algebra $\hat{\mathfrak{g}}$. Then $\hat{G}$ acquires a $GL(n, \mathbb{C})$ -structure, obtained by complexifying (23) and then embedding the complex matrix in $GL(4n, \mathbb{R})$. The induced hypercomplex structure on $\hat{G}$ comes about as a result of inclusions

$$GL(n, \mathbb{C}) \subset GL(n, \mathbb{H}) \subset GL(4n, \mathbb{R}).$$

Before concluding the G -structure approach, we consider cases in which a complex product structure reduces from $GL(n, \mathbb{R})$ to either (i) the orthogonal group $O(n)$, or (ii) the symplectic group $Sp(n/2, \mathbb{R})$.

(i) $O(n)$ represents the subgroup preserving a scalar product g on T_+ or T_- . These two representations are equivalent by means of J , so we extend g to $T = \mathbb{R}^{2n}$ by the requirement that

$$g(JX, JY) = g(X, Y), \quad g(EX, EY) = g(X, Y),$$

for all tangent vectors X, Y . A 2-form is now defined in the usual way, by setting

$$\omega(X, Y) = g(X, JY).$$

If this is closed on a complex product manifold, then g is a reducible *Kähler metric*, and ∇^{CP} coincides with the Levi-Civita connection ∇^g . It is however possible to drop the integrability assumption on J , and consider the complementary *Lagrangian foliations* defined by the distributions $T_{\pm}$ on a symplectic manifold.

(ii) $Sp(n/2, \mathbb{R})$ represents the group preserving a non-degenerate 2-form ω_1 on T_+ or T_- , each of which must then have even dimension $n = 2m$. We may extend ω_1 to T by the requirement that

$$\omega_1(JX, JY) = \omega(X, Y), \quad \omega_1(EX, EY) = \omega_1(X, Y),$$

in analogy to g above. A symmetric bilinear form h is now defined by setting

$$h(X, Y) = \omega_1(JX, Y),$$

and it follows that $h(EX, EY) = -h(X, Y)$; thus $T_{\pm}$ are isotropic and h has signature $(2m, 2m)$. Additional 2-forms are defined by setting

$$\omega_2(X, Y) = h(X, EY), \quad \omega_3(X, Y) = h(X, JEY).$$

A manifold is called *hypersymplectic* if it has such an $Sp(n/2, \mathbb{R})$ structure for which all three forms ω_i are closed [17]. Such manifolds are also called *neutral hyperkähler* [18, 14], and admit tensors J, E, F satisfying (6). The complex 2-form $\Omega = \omega_2 + i\omega_3$ has type $(2, 0)$ relative to J and the fact that Ω^m is closed and non-zero implies that J is integrable. It follows that h is pseudo-Kähler, and that ∇^{CP} coincides with ∇^h . The subspaces $T_{\pm}$ are Lagrangian relative to ω_2 , though it is more usual to seek submanifolds that are special Lagrangian relative to the forms ω_1 and $\text{Im}(\Omega^m)$ (meaning that these restrict to zero).

In conclusion, a hypersymplectic manifold always has an underlying complex product structure (with n even), in the same sort of way that a hyperkähler manifold has a sub-ordinate hypercomplex structure.

6. COMPLEX PRODUCT STRUCTURES ON 4-DIMENSIONAL LIE ALGEBRAS

In this section we shall consider some concrete examples of 4-dimensional Lie algebras carrying complex product structures. We will determine the associated isomorphisms with bicrossproduct Lie algebras and also the corresponding local double Lie groups.

Let us first make the following observation. If a 4-dimensional Lie algebra admits a complex product structure, then it can be written as the sum of two 2-dimensional subalgebras. We will use the fact that any 2-dimensional Lie algebra is either abelian or isomorphic to $\mathfrak{aff}(\mathbb{R})$ (see Example 2.8 (ii)).

We will consider separately the solvable Lie algebras and the non-solvable Lie algebras. It is easy to see that there are only two of the latter, using the Levi decomposition and the classification of 3-dimensional simple Lie algebras. The only possibilities are the reductive Lie algebras $\mathbb{R} \oplus \mathfrak{so}(3)$ and $\mathbb{R} \oplus \mathfrak{sl}(2, \mathbb{R}) \cong \mathfrak{gl}(2, \mathbb{R})$.

6.1. Reductive case.

Example 6.1. Let us begin with $\mathbb{R} \oplus \mathfrak{sl}(2, \mathbb{R}) \cong \mathfrak{gl}(2, \mathbb{R})$. This Lie algebra has a basis $\{W, X, Y, Z\}$ with Z central and non-zero brackets given by

$$[W, X] = 2X, \quad [W, Y] = -2Y, \quad [X, Y] = W.$$

We already know from Proposition 3.5 that $\mathfrak{gl}(2, \mathbb{R})$ carries a complex product structure $\{J, E\}$ given by right multiplication with the standard almost complex and almost product structures J_0 and E_0 on $\mathbb{R}^{2n}$. Identifying W, X, Y, Z with (2×2) matrices, it is easy to see that

$$JW = -(X + Y), \quad JX = \frac{1}{2}(W + Z), \quad JY = \frac{1}{2}(W - Z), \quad JZ = -X + Y,$$

while

$$EW = Z, \quad EX = -X, \quad EY = Y, \quad EZ = W.$$

The eigenspaces for E are the following subalgebras of $\mathfrak{gl}(2, \mathbb{R})$:

$$\mathfrak{g}_+ = \text{span}\{Y, W + Z\}, \quad \mathfrak{g}_- = \text{span}\{X, W - Z\}.$$

Observe that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are isomorphic to $\mathfrak{gl}(2, \mathbb{R})$; therefore the associated double Lie algebra is $(\mathfrak{gl}(2, \mathbb{R}), \mathfrak{aff}(\mathbb{R}), \mathfrak{aff}(\mathbb{R}))$ and its local double Lie group is $(GL(2, \mathbb{R}), G_+, G_-)$, where both G_+ and G_- are locally isomorphic to $Aff(\mathbb{R})$.

Remark. Sasaki classified the complex structures on $\mathfrak{gl}(2, \mathbb{R})$ in [28]. They are parametrized by $d \in \mathbb{C} \setminus \{0\}$ on the curve $\text{Re}(1/d) = -1$. The complex structure which appears above corresponds to the case $d = -1$.

Now consider the other 4-dimensional reductive Lie algebra.

Proposition 6.2. *The Lie algebra $\mathbb{R} \oplus \mathfrak{so}(3)$ does not support any complex product structure.*

Proof. The compact Lie algebra $\mathfrak{g} = \mathbb{R} \oplus \mathfrak{so}(3)$ admits an invariant inner product $\langle \cdot, \cdot \rangle$. Consider a 2-dimensional Lie subalgebra $\mathfrak{u}$ of $\mathfrak{g}$. As $\mathfrak{u}$ is a 2-dimensional Lie algebra, we know that $\mathfrak{u}$ is abelian or isomorphic to $\mathfrak{aff}(\mathbb{R})$. If $\mathfrak{u} \cong \mathfrak{aff}(\mathbb{R})$, there is a basis $\{U_1, U_2\}$ of $\mathfrak{u}$ with $[U_1, U_2] = U_2$. Thus

$$\langle [U_1, U_2], U_2 \rangle = \langle U_2, U_2 \rangle = \|U_2\|^2,$$

but, using the invariance of $\langle \cdot, \cdot \rangle$,

$$\langle [U_1, U_2], U_2 \rangle = -\langle U_2, [U_1, U_2] \rangle,$$

from where we obtain $\langle [U_1, U_2], U_2 \rangle = 0$. Hence, $\|U_2\|^2 = 0$, a contradiction. Therefore, $\mathfrak{u}$ must be abelian.

Let us now suppose that $\mathfrak{g}$ admits a complex product structure $\{J, E\}$ with associated double Lie algebra $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$. As we have just seen, both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are abelian. For $U, U' \in \mathfrak{g}_+$ and $V, V' \in \mathfrak{g}_-$, we compute

$$\langle [U, V], U' \rangle = -\langle V, [U, U'] \rangle = 0$$

and

$$\langle [U, V], V' \rangle = -\langle [V, U], V' \rangle = -\langle U, [V, V'] \rangle = 0$$

because $\mathfrak{g}_\pm$ are abelian. Hence $[U, V] = 0$ and then $\mathfrak{g}$ should be abelian, which is false. The proof is now complete. $\square$

6.2. Solvable case.

Example 6.3. Let $\mathfrak{g}$ be the Lie algebra defined by $\mathfrak{g} = \text{span}\{A, B, C, D\}$ with non-zero Lie bracket relations given by

$$[A, B] = B, [A, C] = -C, [A, D] = -D.$$

$\mathfrak{g}$ lies in the class A2 of the classification made in [26], with $\lambda = -1$. It follows that $\mathfrak{g}$ admits only one complex structure J , up to isomorphism, given by

$$JA = B, JC = D.$$

Consider the following subalgebras of $\mathfrak{g}$:

$$\mathfrak{g}_+ = \text{span}\{A - D, C\}, \quad \mathfrak{g}_- = \text{span}\{B + C, D\}.$$

Set $E|_{\mathfrak{g}_+} = \mathbf{1}$, $E|_{\mathfrak{g}_-} = -\mathbf{1}$; the endomorphism E of $\mathfrak{g}$ is clearly a product structure on $\mathfrak{g}$ which anticommutes with the complex structure J , giving rise then to a complex product structure on $\mathfrak{g}$. Thus, there exists an LSA structure on $\mathfrak{g}_+$ and $\mathfrak{g}_-$, which can be extended to a bilinear product on $\mathfrak{g}$. We already know that this product satisfies condition (10) but we will show that it does not satisfy (9). Denote by L_v the endomorphism of $\mathfrak{g}$ given by left-multiplication with $v \in \mathfrak{g}$. Using equation (20), we obtain

$$\begin{aligned} L_{A-D}(A-D) &= A-D, & L_{B+C}(A-D) &= 2C, \\ L_{A-D}(C) &= -C, & L_{B+C}(C) &= 0, \\ L_{A-D}(B+C) &= B+C, & L_{B+C}(B+C) &= 2D, \\ L_{A-D}(D) &= -D, & L_{B+C}(D) &= 0. \end{aligned}$$

and $L_C \equiv 0$, $L_D \equiv 0$. Now, setting $x = A - D$, $y = B + C$, $z = A - D$ in (9), we get

$$x \cdot (y \cdot z) - (x \cdot y) \cdot z = -4C,$$

while

$$y \cdot (x \cdot z) - (y \cdot x) \cdot z = 2C,$$

and hence (9) does not hold.

Example 6.4. Consider the Lie algebra defined by $\mathfrak{g} = \text{span}\{X, Y, Z, W\}$ with the only non-zero bracket given by $[X, Y] = Z$. This Lie algebra is isomorphic to $\mathfrak{h}_3 \oplus \mathbb{R}$, a direct sum of ideals, where $\mathfrak{h}_3$ denotes the 3-dimensional Heisenberg Lie algebra. This Lie algebra belongs to the class S1 in the classification made in [30], from where it follows that there is, up to equivalence, only one complex structure on $\mathfrak{g}$ and it is given by

$$JX = Y, \quad JZ = W.$$

Proposition 6.5. $\mathfrak{h}_3 \oplus \mathbb{R}$ admits complex product structures. With each of these structures, the associated double Lie algebra is $(\mathfrak{h}_3 \oplus \mathbb{R}, \mathbb{R}^2, \mathbb{R}^2)$ and the associated local double Lie group is (G, G_+, G_-) , where G is locally isomorphic to $H_3 \times \mathbb{R}$ and G_+, G_- are locally isomorphic to $\mathbb{R}^2$.

Proof. It can be shown that the product structures on $\mathfrak{g}$ which anticommute with J are parametrized by

$$E_\theta = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & \cos \theta & \sin \theta \\ 0 & 0 & \sin \theta & -\cos \theta \end{pmatrix},$$

where the ordered basis used for the matrix representation is $\{X, Y, Z, W\}$. The pairs $\{J, E_\theta\}$ with $\theta \in [0, 2\pi)$ exhaust all the complex product structures on $\mathfrak{g}$, up to equivalence.

For a fixed $\theta \in [0, 2\pi)$, let us establish the decomposition of $\mathfrak{g}$ into a sum of subalgebras. In order to do so, we simply have to determine the eigenspaces of E_θ . The eigenspace $\mathfrak{g}_+$ corresponding to the eigenvalue $+1$ is given by

$$\mathfrak{g}_+ = \text{span}\{X, \cos_{\theta/2} Z + \sin_{\theta/2} W\},$$

while the eigenspace $\mathfrak{g}_-$ corresponding to the eigenvalue -1 is given by

$$\mathfrak{g}_- = \text{span}\{Y, -\sin_{\theta/2} Z + \cos_{\theta/2} W\}.$$

(We indicate the trigonometric arguments as subscripts for visual clarity.) Observe that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are abelian Lie algebras. Hence, we have $\mathfrak{g} = \mathbb{R}^2 \ltimes \mathbb{R}^2$ for suitable representations and certain LSA structures on each $\mathbb{R}^2$. $\square$

Example 6.6. Let $\mathfrak{g} = \text{span}\{A, B, C, D\}$ be the Lie algebra with non-zero Lie bracket relations given by

$$[A, B] = B, \quad [A, C] = C, \quad [A, D] = D.$$

Then $\mathfrak{g}$ lies in the class A4 of the classification [26], and is the Lie algebra corresponding to the real hyperbolic space $\mathbb{R}H^4$, i.e. the simply connected Lie group S with $L(S) = \mathfrak{g}$ acts simply transitively on $\mathbb{R}H^4$. It follows that $\mathfrak{g}$ admits only one complex structure J , up to isomorphism, given by

$$JA = B, \quad JC = D.$$

Proposition 6.7. *$\mathfrak{g}$ admits complex product structures. The possible associated double Lie algebras are either $(\mathfrak{g}, \mathfrak{aff}(\mathbb{R}), \mathfrak{aff}(\mathbb{R}))$ or $(\mathfrak{g}, \mathbb{R}^2, \mathfrak{aff}(\mathbb{R}))$ and the associated local double Lie groups are (S, G_+, G_-) , where both G_+, G_- are both locally isomorphic to $\text{Aff}(\mathbb{R})$, or G_+ is locally isomorphic to $\mathbb{R}^2$ and G_- to $\text{Aff}(\mathbb{R})$.*

Proof. The following are examples of product structures on $\mathfrak{g}$ which anticommute with the complex structure J :

$$\begin{aligned} E_\theta &= \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ \sin \theta & -\cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, & E'_\theta &= \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ \sin \theta & -\cos \theta & 0 & 0 \\ -\sin \theta & 1 + \cos \theta & 1 & 0 \\ 1 + \cos \theta & \sin \theta & 0 & -1 \end{pmatrix}, \\ E''_\theta &= \begin{pmatrix} \cos \theta & \sin \theta & -\sin \theta & 1 + \cos \theta \\ \sin \theta & -\cos \theta & 1 + \cos \theta & \sin \theta \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, & \tilde{E} &= \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -2 & 0 & 1 & 0 \\ 0 & 2 & 0 & -1 \end{pmatrix}. \end{aligned}$$

The ordered basis used for the matrix representation is $\{A, B, C, D\}$. It can be shown that every complex product structure on $\mathfrak{g}$ is equivalent to $\{J, E\}$ where J is as above and

$E \in \{E_\theta : \theta \in [0, 2\pi)\} \cup \{E'_\theta : \theta \in [0, 2\pi), \theta \neq \pi\} \cup \{E''_\theta : \theta \in [0, 2\pi), \theta \neq \pi\} \cup \{\tilde{E}\}$, and these are all inequivalent.

For $\theta \in [0, 2\pi)$, let us compute the eigenspaces corresponding to E_θ . The eigenspace $\mathfrak{g}_+$ corresponding to the eigenvalue $+1$ is given by

$$\mathfrak{g}_+ = \text{span}\{C, \cos_{\theta/2} A + \sin_{\theta/2} B\},$$

while the eigenspace $\mathfrak{g}_-$ corresponding to the eigenvalue -1 is

$$\mathfrak{g}_- = \text{span}\{D, -\sin_{\theta/2} A + \cos_{\theta/2} B\}.$$

Observe that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are isomorphic to $\mathfrak{aff}(\mathbb{R})$, hence we have an isomorphism $\mathfrak{g} \cong \mathfrak{aff}(\mathbb{R}) \ltimes \mathfrak{aff}(\mathbb{R})$, for suitable representations and certain LSA structures on each $\mathfrak{aff}(\mathbb{R})$.

For $\theta \in [0, 2\pi)$, $\theta \neq \pi$, let us consider the complex product structures $\{J, E'_\theta\}$. The eigenspaces of E'_θ are

$$\begin{aligned} \mathfrak{g}'_+ &= \text{span}\{C, \cos_{\theta/2} A + \sin_{\theta/2} B + \cos_{\theta/2} D\}, \\ \mathfrak{g}'_- &= \text{span}\{D, -\sin_{\theta/2} A + \cos_{\theta/2} B - \cos_{\theta/2} C\}. \end{aligned}$$

Observe that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are isomorphic to $\mathfrak{aff}(\mathbb{R})$, hence $\mathfrak{g} \cong \mathfrak{aff}(\mathbb{R}) \ltimes \mathfrak{aff}(\mathbb{R})$, for suitable representations and certain LSA structures on each $\mathfrak{aff}(\mathbb{R})$.

For $\theta \in [0, 2\pi)$, $\theta \neq \pi$, let us consider the complex product structures $\{J, E''_\theta\}$. The eigenspaces of E''_θ are

$$\begin{aligned} \mathfrak{g}''_+ &= \text{span}\{\cos_{\theta/2} A + \sin_{\theta/2} B, -\cos_{\theta/2} A + \sin_{\theta/2} C\}, \\ \mathfrak{g}''_- &= \text{span}\{-\sin_{\theta/2} A + \cos_{\theta/2} B, -\cos_{\theta/2} B + \sin_{\theta/2} D\}. \end{aligned}$$

Note that both $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are isomorphic to $\mathfrak{aff}(\mathbb{R})$, hence $\mathfrak{g} \cong \mathfrak{aff}(\mathbb{R}) \ltimes \mathfrak{aff}(\mathbb{R})$, for suitable representations and certain LSA structures on each $\mathfrak{aff}(\mathbb{R})$.

Finally, the eigenspaces for $\tilde{E}$ are

$$\tilde{\mathfrak{g}}_+ = \text{span}\{C, B + D\}, \quad \tilde{\mathfrak{g}}_- = \text{span}\{D, A + C\}.$$

Note that $\tilde{\mathfrak{g}}_+$ is abelian while $\tilde{\mathfrak{g}}_-$ is isomorphic to $\mathfrak{aff}(\mathbb{R})$. Thus, in this case we have an isomorphism $\mathfrak{g} \cong \mathbb{R}^2 \ltimes \mathfrak{aff}(\mathbb{R})$, for suitable representations and certain LSA structures on each summand. $\square$

In contrast to above, not every solvable Lie algebra (among those carrying a complex structure) admits a complex product structure.

Example 6.8. Let $\mathfrak{g}$ be the Lie algebra given by $\mathfrak{g} = \text{span}\{A, X, Y, Z\}$ with Lie bracket relations

$$[X, Y] = Z, \quad [A, X] = -Y, \quad [A, Y] = X.$$

This Lie algebra belongs to the class *H2* of the classification made in [26] and it follows that it admits, up to equivalence, only one complex structure J , which is given by

$$JA = Z, \quad JX = Y.$$

We will show that this Lie algebra does not admit any complex product structure, using

Lemma 6.9. *$\mathfrak{g}$ does not contain a non-abelian 2-dimensional Lie subalgebra.*

Proof. Suppose that $\mathfrak{u}$ is a non-abelian 2-dimensional Lie subalgebra of $\mathfrak{g}$. Then it is isomorphic to $\mathfrak{aff}(\mathbb{R})$ and hence there is a basis $\{U, V\}$ of $\mathfrak{u}$ such that $[U, V] = V$. If $U = a_1A + x_1X + y_1Y + z_1Z$, $V = a_2A + x_2X + y_2Y + z_2Z$, the relation $[U, V] = V$ implies at once that $a_2 = 0$ and

$$\begin{cases} a_1y_2 = x_2, \\ a_1x_2 = -y_2, \\ x_1y_2 - y_1x_2 = z_2. \end{cases}$$

From the first two equations we obtain that $y_2(a_1^2 + 1) = 0$. Since we are working over $\mathbb{R}$, we have that $y_2 = 0$. But this implies that $x_2 = 0$ and hence $z_2 = 0$, showing that $U = 0$, which is a contradiction. $\square$

Let us suppose now that $\mathfrak{g}$ admits a complex product structure. Then, we have $\mathfrak{g} = \mathfrak{g}_+ \oplus \mathfrak{g}_-$ with $\mathfrak{g}_+$ and $\mathfrak{g}_- = J\mathfrak{g}_+$ subalgebras of $\mathfrak{g}$. From the lemma, we obtain that both $\mathfrak{g}_\pm$ are abelian. It can be seen that in this case the complex structure J satisfies the condition

$$(24) \quad [JX, JY] = [X, Y]$$

for all $X, Y \in \mathfrak{g}$. A complex structure J on a Lie algebra $\mathfrak{g}$ satisfying (24) is called *abelian*, and only solvable Lie algebras admit such structures [13]. A result in [6] states that if a Lie algebra with commutator ideal of codimension 1 admits an abelian complex structure then it is isomorphic to $\mathfrak{aff}(\mathbb{R})$. Since $\mathfrak{g}$ is not isomorphic to $\mathfrak{aff}(\mathbb{R})$, we obtain a contradiction and then $\mathfrak{g}$ cannot carry a complex product structure.

6.3. Induced hypercomplex structures. In this subsection we will give some applications of Theorem 3.3, using the examples studied in §6.1 and §6.2. We will use the following notation: for $X \in \mathfrak{g}$, we denote $\hat{X} = iX$ in $\mathfrak{g}^{\mathbb{C}}$.

Example 6.10. Let us begin with $\mathfrak{gl}(2, \mathbb{R})$. Consider on this Lie algebra the complex product structure given in §6.1 and let $\{\mathbb{I}, \mathbb{J}\}$ be the hypercomplex structure on $\mathfrak{gl}(2, \mathbb{C}) = \widehat{\mathfrak{gl}(2, \mathbb{R})}$ given by Theorem 3.3 and (14). $\mathfrak{gl}(2, \mathbb{C})$ is an 8-dimensional real Lie algebra with a basis $\{W, X, Y, Z, \hat{W}, \hat{X}, \hat{Y}, \hat{Z}\}$ and Lie bracket given by

$$\begin{aligned} [W, X] &= -[\hat{W}, \hat{X}] = 2X, & [W, Y] &= -[\hat{W}, \hat{Y}] = -2Y, & [X, Y] &= -[\hat{X}, \hat{Y}] = W, \\ [W, \hat{X}] &= [\hat{W}, X] = 2\hat{X}, & [W, \hat{Y}] &= -[\hat{W}, Y] = -2\hat{Y}, & [\hat{X}, Y] &= -[X, \hat{Y}] = \hat{W}, \end{aligned}$$

The hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ is given by

$$\begin{aligned} \mathbb{I}W &= \hat{Z}, & \mathbb{I}X &= -\hat{X}, & \mathbb{I}Y &= \hat{Y}, & \mathbb{I}Z &= \hat{W}, \\ \mathbb{I}\hat{W} &= -Z, & \mathbb{I}\hat{X} &= -X, & \mathbb{I}\hat{Y} &= Y, & \mathbb{I}\hat{Z} &= -W, \end{aligned}$$

and

$$\begin{aligned} \mathbb{J}W &= -(X + Y), & \mathbb{J}X &= \frac{1}{2}(W + Z), & \mathbb{J}Y &= \frac{1}{2}(W - Z), & \mathbb{J}Z &= -X + Y, \\ \mathbb{J}\hat{W} &= -(\hat{X} + \hat{Y}), & \mathbb{J}\hat{X} &= \frac{1}{2}(\hat{W} + \hat{Z}), & \mathbb{J}\hat{Y} &= \frac{1}{2}(\hat{W} - \hat{Z}), & \mathbb{J}\hat{Z} &= -\hat{X} + \hat{Y}. \end{aligned}$$

Example 6.11. We continue here Example 6.3. $\hat{\mathfrak{g}} = (\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$ is an 8-dimensional solvable Lie algebra with a basis $\{A, B, C, D, \hat{A}, \hat{B}, \hat{C}, \hat{D}\}$ and non-zero Lie brackets given by

$$\begin{aligned} [A, B] &= -[\hat{A}, \hat{B}] = B, & [A, C] &= -[\hat{A}, \hat{C}] = -C, & [A, D] &= -[\hat{A}, \hat{D}] = -D, \\ [A, \hat{B}] &= [\hat{A}, B] = \hat{B}, & [A, \hat{C}] &= [\hat{A}, C] = -\hat{C}, & [A, \hat{D}] &= [\hat{A}, D] = -\hat{D}. \end{aligned}$$

The complex product structure considered on $\mathfrak{g}$ gives rise to a hypercomplex structure $\{\mathbb{I}, \mathbb{J}\}$ on $\hat{\mathfrak{g}}$. This hypercomplex structure is given by

$$\begin{aligned} \mathbb{I}A &= \hat{A} - 2\hat{D}, & \mathbb{I}B &= -\hat{B} - 2\hat{C}, & \mathbb{I}C &= \hat{C}, & \mathbb{I}D &= -\hat{D}, \\ \mathbb{I}\hat{A} &= -A + 2D, & \mathbb{I}\hat{B} &= B + 2C, & \mathbb{I}\hat{C} &= -C, & \mathbb{I}\hat{D} &= D, \end{aligned}$$

and

$$\mathbb{J}A = B, \quad \mathbb{J}C = D, \quad \mathbb{J}\hat{A} = \hat{B}, \quad \mathbb{J}\hat{C} = \hat{D}.$$

According to Example 6.3 and Corollary 5.3 we obtain that the Obata connection on $\hat{\mathfrak{g}}$ (or on any Lie group G with $L(G) = \hat{\mathfrak{g}}$) is not flat.

Example 6.12. We continue here Example 6.4. If $\mathfrak{g} = \mathfrak{h}_3 \oplus \mathbb{R}$, then $\hat{\mathfrak{g}} = (\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$ is an 8-dimensional 2-step nilpotent Lie algebra with a basis $\{X, Y, Z, W, \hat{X}, \hat{Y}, \hat{Z}, \hat{W}\}$ and non-zero Lie brackets given by

$$[X, Y] = -[\hat{X}, \hat{Y}] = Z, \quad [X, \hat{Y}] = [\hat{X}, Y] = \hat{Z}.$$

The hypercomplex structure $\{\mathbb{I}_{\theta}, \mathbb{J}\}$ on $\hat{\mathfrak{g}}$ given by Theorem 3.3 is

$$\begin{aligned} \mathbb{I}_{\theta}X &= \hat{X}, & \mathbb{I}_{\theta}Y &= -\hat{Y}, & \mathbb{I}_{\theta}Z &= \cos_{\theta} \hat{Z} + \sin_{\theta} \hat{W}, & \mathbb{I}_{\theta}W &= \sin_{\theta} \hat{Z} - \cos_{\theta} \hat{W}, \\ \mathbb{I}_{\theta}\hat{X} &= -X, & \mathbb{I}_{\theta}\hat{Y} &= Y, & \mathbb{I}_{\theta}\hat{Z} &= -\cos_{\theta} Z - \sin_{\theta} W, & \mathbb{I}_{\theta}\hat{W} &= -\sin_{\theta} Z + \cos_{\theta} W, \\ \mathbb{J}X &= Y, & \mathbb{J}Z &= W, & \mathbb{J}\hat{X} &= \hat{Y}, & \mathbb{J}\hat{Z} &= \hat{W}. \end{aligned}$$

Remark. A classification of 8-dimensional nilpotent Lie groups carrying an abelian hypercomplex structure (meaning all its complex structures satisfy (24)) was given in [12]. According to this classification, if G is a Lie group whose Lie algebra is the one considered in the previous example, then G is isomorphic to a trivial extension of a group of type H with centre of dimension 2.

Example 6.13. We continue here Example 6.6. $\hat{\mathfrak{g}} = (\mathfrak{g}^{\mathbb{C}})_{\mathbb{R}}$ is an 8-dimensional solvable Lie algebra with a basis $\{A, B, C, D, \hat{A}, \hat{B}, \hat{C}, \hat{D}\}$ and non-zero Lie brackets given by

$$\begin{aligned} [A, B] &= -[\hat{A}, \hat{B}] = B, & [A, C] &= -[\hat{A}, \hat{C}] = C, & [A, D] &= -[\hat{A}, \hat{D}] = D, \\ [A, \hat{B}] &= [\hat{A}, B] = \hat{B}, & [A, \hat{C}] &= [\hat{A}, C] = \hat{C}, & [A, \hat{D}] &= [\hat{A}, D] = \hat{D}. \end{aligned}$$

Each complex product structure on $\mathfrak{g}$ gives rise to a hypercomplex structure on $\hat{\mathfrak{g}}$.

If we take $\{J, E_{\theta}\}$, we have the hypercomplex structure $\{\mathbb{I}_{\theta}, \mathbb{J}\}$, where

$$\begin{aligned} \mathbb{I}_{\theta}A &= \cos_{\theta} \hat{A} + \sin_{\theta} \hat{B}, & \mathbb{I}_{\theta}B &= \sin_{\theta} \hat{A} - \cos_{\theta} \hat{B}, & \mathbb{I}_{\theta}C &= \hat{C}, & \mathbb{I}_{\theta}D &= -\hat{D}, \\ \mathbb{I}_{\theta}\hat{A} &= -\cos_{\theta} A - \sin_{\theta} B, & \mathbb{I}_{\theta}\hat{B} &= -\sin_{\theta} A + \cos_{\theta} B, & \mathbb{I}_{\theta}\hat{C} &= -C, & \mathbb{I}_{\theta}\hat{D} &= D, \end{aligned}$$

and

$$\mathbb{J}A = B, \quad \mathbb{J}C = D, \quad \mathbb{J}\hat{A} = \hat{B}, \quad \mathbb{J}\hat{C} = \hat{D}.$$

Considering $\{J, E'_\theta\}$, we have the hypercomplex structure $\{\mathbb{I}'_\theta, \mathbb{J}\}$, with $\mathbb{J}$ as above and

$$\begin{aligned}\mathbb{I}'_\theta A &= \cos_\theta \hat{A} + \sin_\theta \hat{B} - \sin_\theta \hat{C} + (1 + \cos_\theta) \hat{D}, \\ \mathbb{I}'_\theta B &= \sin_\theta \hat{A} - \cos_\theta \hat{B} + (1 + \cos_\theta) \hat{C} + \sin_\theta \hat{D}, \\ \mathbb{I}'_\theta C &= \hat{C}, \quad \mathbb{I}'_\theta D = -\hat{D}, \\ \mathbb{I}'_\theta \hat{A} &= -\cos_\theta A - \sin_\theta B + \sin_\theta C - (1 + \cos_\theta) D, \\ \mathbb{I}'_\theta \hat{B} &= -\sin_\theta A + \cos_\theta B - (1 + \cos_\theta) C - \sin_\theta D, \\ \mathbb{I}'_\theta \hat{C} &= -C, \quad \mathbb{I}'_\theta \hat{D} = D.\end{aligned}$$

For $\{J, E''_\theta\}$, we have the hypercomplex structure $\{\mathbb{I}''_\theta, \mathbb{J}\}$, where $\mathbb{J}$ is as above and

$$\begin{aligned}\mathbb{I}''_\theta A &= \cos_\theta \hat{A} + \sin_\theta \hat{B}, \quad \mathbb{I}''_\theta B = \sin_\theta \hat{A} - \cos_\theta \hat{B}, \\ \mathbb{I}''_\theta C &= -\sin_\theta \hat{A} + (1 + \cos_\theta) \hat{B} + \hat{C}, \quad \mathbb{I}''_\theta D = (1 + \cos_\theta) \hat{A} + \sin_\theta \hat{B} - \hat{D}, \\ \mathbb{I}''_\theta \hat{A} &= -\cos_\theta A - \sin_\theta B, \quad \mathbb{I}''_\theta \hat{B} = -\sin_\theta A + \cos_\theta B, \\ \mathbb{I}''_\theta \hat{C} &= \sin_\theta A - (1 + \cos_\theta) B - C, \quad \mathbb{I}''_\theta \hat{D} = -(1 + \cos_\theta) A - \sin_\theta B + D,\end{aligned}$$

Finally, for the complex product structure $\{J, \tilde{E}\}$ we have the hypercomplex structure $\{\tilde{\mathbb{I}}, \mathbb{J}\}$, with $\mathbb{J}$ as above and

$$\begin{aligned}\tilde{\mathbb{I}} A &= -\hat{A} - 2\hat{C}, \quad \tilde{\mathbb{I}} B = \hat{B} + 2\hat{D}, \quad \tilde{\mathbb{I}} C = \hat{C}, \quad \tilde{\mathbb{I}} D = -\hat{D}, \\ \tilde{\mathbb{I}} \hat{A} &= A + 2C, \quad \tilde{\mathbb{I}} \hat{B} = -B - 2D, \quad \tilde{\mathbb{I}} \hat{C} = -C, \quad \tilde{\mathbb{I}} \hat{D} = D.\end{aligned}$$

REFERENCES

- [1] A. Andrada, “Complex product structures on differentiable manifolds”, preprint.
- [2] L. Auslander and L. Markus, “Holonomy of flat affinely connected manifolds”, *Ann. of Math.* **62**, 139–151 (1955).
- [3] T. N. Bailey and M. G. Eastwood, “Complex paraconformal manifolds – their differential geometry and twistor theory”, *Forum Math.* **3**, 61–103 (1991).
- [4] M. L. Barberis, “Hypercomplex structures on four-dimensional Lie groups”, *Proc. Amer. Math. Soc.* **125**, 1043–1054 (1997).
- [5] M. L. Barberis, “Affine connections on homogeneous hypercomplex manifolds”, *J. Geom. Phys.* **32**, 1–13 (1999).
- [6] M. L. Barberis and I. Dotti, “Abelian complex structures on solvable Lie algebras”, to appear in *J. Lie Theory*.
- [7] Y. Benoist, “Une nilvariété non-affine”, *J. Differ. Geom.* **41**, 21–52 (1995).
- [8] N. B. Boyom, “Variétés symplectiques affines”, *Manuscripta Math.* **64**, 1–33 (1989).
- [9] D. Burde, “Left-symmetric structures on simple modular Lie algebras”, *J. Algebra* **169**, 112–138 (1994).
- [10] D. Burde, “Left-invariant affine structures on reductive Lie groups”, *J. Algebra* **181**, 884–902 (1996).
- [11] A. Diatta and A. Medina, “Classical Yang-Baxter equation and left-invariant affine geometry on Lie groups”, preprint 2002: DG/0203198.
- [12] I. Dotti and A. Fino, “Abelian hypercomplex 8-dimensional nilmanifolds”, *Ann. Global Anal. Geom.* **18**, 47–59 (2000).
- [13] I. Dotti and A. Fino, “Hypercomplex nilpotent Lie groups”, in *Global Differential Geometry: The Mathematical Legacy of Alfred Gray*, 310–314, *Contemp. Math.* **241**, Amer. Math. Soc., 2001.
- [14] A. Fino, H. Pedersen, Y. S. Poon, and M. Sørensen, “Neutral Calabi-Yau structures on Kodaira manifolds”, preprint, 2002: <http://www.imada.sdu.dk>

- [15] E. García-Río, Y. Matsushita and R. Vázquez-Lorenzo, “Paraquaternionic Kähler manifolds”, *Rocky Mountain J. Math.* **31**, 237–261 (2001)
- [16] J. Helmstetter, “Radical d’une algèbre symétrique à gauche”, *Ann. Inst. Fourier (Grenoble)* **29**, 17–35 (1979).
- [17] N. Hitchin, “Hypersymplectic quotients”, *Acta Acad. Sci. Tauriensis* **124** suppl, 169–180 (1990).
- [18] H. Kamada, “Neutral hyperKähler structures on primary Kodaira surfaces”, *Tsukuba J. Math.* **23**, 321–332 (1999).
- [19] S. Kaneyuni and M. Kozai, “Paracomplex structures and affine symmetric spaces”, *Tokyo J. Math.* **8**, 81–98 (1985).
- [20] P. Libermann, “Sur les structures presque paracomplexes”, *C. R. Acad. Sci. Paris* **234**, 2517–2519 (1952).
- [21] J.-H. Lu and A. Weinstein, “Poisson Lie groups, dressing transformations and Bruhat decompositions”, *J. Diff. Geom.* **31**, 501–526 (1990).
- [22] S. Majid, “Matched pairs of Lie groups associated to solutions of the Yang-Baxter equations”, *Pacific J. Math.* **141**, 311–332 (1990)
- [23] A. Masuoka, “Extensions of Hopf algebras and Lie bialgebras”, *Trans. Amer. Math. Soc.* **352**, 3837–3879 (2000)
- [24] J. Milnor, “On fundamental groups of complete affinely flat manifolds”, *Adv. in Math.* **25**, 178–187 (1977).
- [25] M. Obata, “Affine connections on manifolds with almost complex, quaternion or hermitian structure”, *Japan. J. Math.* **26** 43–77 (1956).
- [26] G. Ovando, “Invariant complex structures on solvable real Lie groups”, *Manuscripta math.* **103**, 19–30 (2000).
- [27] S. M. Salamon, “Differential geometry of quaternionic manifolds”, *Ann. Sci. École Norm. Sup.* **19**, 31–55 (1986).
- [28] T. Sasaki, “Classification of left invariant complex structures on $GL(2, \mathbb{R})$ and $U(2)$ ”, *Kumamoto J. Sci. (Math.)* **14**, 115–123 (1981).
- [29] D. Segal, “Free left-symmetric algebras and an analogue of the Poincaré-Birkhoff-Witt Theorem”, *J. Algebra* **164** 750–772 (1994).
- [30] J. E. Snow, “Invariant complex structures on four-dimensional solvable real Lie groups”, *manuscripta math.* **66**, 397–412 (1990).
- [31] S. Sternberg, “Lectures on differential geometry”, Prentice-Hall, 1964.

CIEM, FaMAF, Universidad Nacional de Córdoba, Ciudad Universitaria, (5000) Córdoba, Argentina
 andrada@mate.uncor.edu

Dipartimento di Matematica, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy
 salamon@calvino.polito.it